

©Copyright 2016

Christine L. Heumann

Comparison of In-Person Versus Telephone Interviews for Early Syphilis and HIV  
Partner Services in King County, Washington (2010-2014)

Christine L. Heumann

A thesis

submitted in partial fulfillment of the  
requirements for the degree of

Master of Public Health

University of Washington

2016

Committee:

Matthew R. Golden

Lisa E. Manhart

Program Authorized to Offer Degree:

Epidemiology

University of Washington

**Abstract**

Comparison of In-Person Versus Telephone Interviews for Early Syphilis and HIV Partner Services in King County, Washington (2010-2014)

Christine L. Heumann

Chair of the Supervisory Committee:  
Matthew R. Golden, Adjunct Professor  
Epidemiology

**Background:** Early syphilis (ES) and HIV partner services (PS) traditionally use in-person interviews. Telephone interviews may be more efficient, but their relative effectiveness is uncertain. **Methods:** We used multivariable Poisson regression to compare partners named, notified, tested, diagnosed, and treated (ES only) per original patient (OP) who underwent in-person versus telephone PS interview in King County, Washington from 2010-2014. **Results:** For ES (682 in-person, 646 telephone OP interviews), in-person interviews were associated with more partners named (aRR=1.68[95% CI: 1.55-1.82]), notified (aRR=1.39[1.24-1.56]), tested (aRR=1.34[1.16-1.54]), and empirically treated (aRR=1.19[1.03-1.37]) but no difference in infected partners treated (aRR=0.93[0.72-1.21]). For HIV (358 in-person, 489 telephone OP interviews), in-person interviews were associated with more partners named (aRR=1.38[1.18-1.62]), notified (aRR=1.24[1.03-1.50]), and newly diagnosed with HIV (aRR=2.17[1.04-4.50]) but no difference in partners tested (aRR=1.15[0.88-1.52]). **Conclusions:** These data support prioritizing in-person HIV PS and suggest that in-person syphilis PS may not have major benefit over telephone PS.

## **Introduction:**

Public health partner services (PS) seek to identify and notify the sex partners of persons with sexually transmitted diseases (STDs), including human immunodeficiency virus (HIV), with a goal of identifying undiagnosed persons, linking them to care, and assuring their treatment. Public health staff providing PS have traditionally conducted in-person interviews with infected persons, also known as original patients (OPs), to elicit the names of potentially exposed partners. The reason for using in-person meetings for PS is multifactorial, including confidentiality concerns<sup>1</sup>, facilitating collection of specimens for testing<sup>2</sup>, providing treatment and linkage to medical care as appropriate<sup>2</sup>, and the commonly held belief that in-person interviews are more effective.

The most recent Center for Disease Control (CDC) guidelines for STD/HIV partner services recommend that interviews should be conducted in person, but if no reasonable alternative exists, telephone interviews may be conducted.<sup>3</sup> In King County, Washington, disease intervention specialists (DIS) routinely interview persons with HIV and syphilis over the telephone with safeguards in place to ensure privacy and confidentiality. For the most part, DIS only perform in-person interviews if a patient is in the Public Health - Seattle and King County (PHSKC) STD Clinic or at the nearby affiliated hospital, if efforts to interview the OP over the telephone are unsuccessful, or if a person with newly diagnosed HIV does not successfully link to medical care. In-person interviews are commonly performed for patients at the STD clinic or the nearby affiliated hospital because DIS work onsite at the STD clinic. Other health departments have reported using telephone interviews for PS interviews, and some use email and text messaging for PS communications.<sup>4-8</sup>

Given resource constraints, it is important to maximize efficiencies in STD partner services.<sup>9</sup> Telephone interviews may be more efficient than in-person interviews; however, their relative effectiveness is uncertain. A survey of attendees of STD and community clinics as well as a community based organization about hypothetical PS found that there was no significant difference in favorability of in-clinic notification compared to telephone notification from the perspective of OPs or partners.<sup>10</sup> Another study conducted in New York City found that a change in program policy that allowed DIS to conduct telephone interviews of partners exposed to HIV resulted in an increase in number of partners notified and tested, though the proportion of notified partners known to have tested declined.<sup>11</sup> To our knowledge, there have been no prior studies that have compared the outcomes of in-person versus telephone PS interviews for OPs.

In this study, we compared in-person versus telephone PS interviews performed for OPs with early syphilis or new HIV diagnoses from 2010-2014 in King County, Washington to determine if there were differences in numbers of partners named, notified, tested, and newly diagnosed with infection. For early syphilis diagnoses, we also evaluated the number of partners treated and number of infected partners treated.

## **Materials and Methods:**

### *Study design and data sources*

We conducted a population-based retrospective cohort study of OPs who underwent PS interviews in King County, Washington from 2010-2014. Data were extracted and de-identified from PS records completed by DIS at PHSKC, the state's

partner services data management system, and the enhanced HIV/AIDS Reporting System (eHARS). The University of Washington Institution Review Board (IRB) defined the project as a program evaluation that did not require human subjects research review and approval.

### *Study population*

The study population included all King County, Washington residents reported to PHSKC with early syphilis or newly diagnosed HIV infection who were interviewed by DIS for PS between 2010 and 2014. We excluded records from OPs with incomplete interviews. We defined cases as early syphilis (primary, secondary, or early latent syphilis) based on the staging evaluation of a single DIS who reviews all syphilis cases for the county. We considered HIV diagnoses to be new if they were never previously reported in Washington State, were not defined to have been previously reported in another state based on CDC de-duplication procedures, and the patient denied a prior HIV diagnosis during their PS interview.

### *Study data and outcomes*

Study data included information obtained through interviews and subsequent investigations with the OP and their partners. The primary exposure of interest for this analysis was OP interview method (telephone or in-person). DIS recorded this information using a checkbox on the PS interview. Covariates related to the OP included age, gender, race/ethnicity, sexual orientation, diagnosing clinic type, time from

diagnosis to interview, HIV status (for ES only), and the DIS who performed the interview.

Data on partner outcomes was extracted from partner PS records and linked to the corresponding OP by case number. The outcome variables created for early syphilis were number of partners named, notified, tested for syphilis, tested for HIV, empirically treated for syphilis, diagnosed with syphilis, diagnosed with HIV, and diagnosed with and treated for syphilis. The outcome variables created for newly diagnosed HIV were number of partners named, notified, tested for HIV, and diagnosed with HIV. Similar to prior PS program evaluations, we assessed each outcome as an index of total numbers of partners divided by total number of OP interviews.<sup>12,13</sup> Each index therefore represents the mean number of partners with the outcome per OP interviewed. The six indices evaluated in this study were 1) contact index: number of partners named per OP interviewed, 2) notification index: number of partners notified per OP interviewed, 3) test index: number of partners tested per OP interviewed, 4) epidemiologic index: number of partners empirically treated per OP interviewed, 5) case-finding index: number of partners newly diagnosed with infection per OP interviewed, 6) brought-to-treatment index: number of infected partners treated per OP interviewed.

### *Statistical Analyses*

We used Stata 14.1 software for all statistical analyses.<sup>14</sup> We first compared key covariates between OPs that underwent in-person interviews and OPs that underwent telephone interviews. Age and time from diagnosis to interview were evaluated as

continuous variables; Wilcoxon rank sum tests were used to evaluate between-group differences in medians. All other variables were categorical and we used a Pearson's chi squared test to evaluate between-group differences. Indices were first calculated at the population level with pooled counts of partner outcomes divided by number of OPs interviewed by interview method. For the syphilis brought-to-treatment index and for the HIV case-finding index, we also calculated verified indices that only included outcomes that were verified by the health department through direct communication with the partner, direct communication with the provider, or medical record review. We used unadjusted Poisson regression models to calculate indices and relative risks (RR) with 95% confidence intervals (CI). We then used multivariable Poisson regression models, which adjusted for OP age, gender, race/ethnicity, sexual orientation, time from OP diagnosis to interview, diagnosing clinic type, and the DIS who performed the interview, to calculate adjusted RRs with 95% CIs. For early syphilis outcomes, we also adjusted for HIV status. Next we evaluated whether in-person interviews were conducted in the field or at the STD clinic and repeated the multivariable Poisson regression models to compare field-based interviews versus telephone interviews and STD clinic interviews versus telephone interviews. Lastly, we evaluated the association between OP age, gender, race/ethnicity, sexual orientation, and time from OP diagnosis to interview and the syphilis brought-to-treatment and HIV case-finding indices.

## **Results:**

### *Early Syphilis*

Clinicians and laboratories reported a total of 1,630 OPs with early syphilis, of whom 1,328 underwent a complete PS interview and were included in the analysis. A total of 682 OPs underwent in-person interviews and 646 OPs underwent telephone interviews (Table 1). Compared to persons interviewed by telephone, OPs interviewed in person were younger, interviewed sooner after syphilis diagnosis, and were more likely to be non-white, HIV negative, and diagnosed in the PHSKC STD Clinic or an affiliated community-based organization (CBO) ( $p < 0.001$ ). The percentage of cases receiving in-person and telephone interviews varied by DIS ( $p < 0.001$ ).

Compared to telephone interviews, in-person interviews resulted in significantly more partners per OP being named, notified, tested for syphilis, tested for HIV, and empirically treated for syphilis, with adjusted relative risks (aRRs) ranging from 1.19 to 1.68 (Table 2). Despite these differences, there was no significant difference between in-person and telephone interviews in syphilis case-finding or the brought-to-treatment indices. Similarly, the HIV case-finding index among persons with early syphilis did not vary significantly between OPs interviewed in-person and by telephone.

Of the 682 in-person interviews, 26 were field investigations, 624 took place in the STD clinic. The remaining 32 in-person interviews did not have a location recorded. We found fewer differences between field-based interviews and telephone interviews with no significant difference in the contact, syphilis test, HIV test, epidemiologic, syphilis case-finding, HIV case-finding, and syphilis brought-to-treatment indices. Field-based interviews were associated with a lower notification index than telephone

interviews (aRR 0.60, 95% CI 0.38-0.98). The evaluation of STD clinic interviews versus telephone interviews resulted in the same findings as our primary analysis. Overall, white race (RR 1.28, 95% CI 1.01-1.62) and a time from diagnosis to interview of less than 14 days (RR 1.44, 95% CI 1.16-1.80) were associated with higher brought-to-treatment indices.

### *Newly diagnosed HIV*

Clinicians and laboratories reported a total of 1,376 in-state OPs with new HIV diagnoses, of whom 847 underwent a complete PS interview and were included in the analysis. A total of 358 OPs underwent in-person interviews and 489 OPs underwent telephone interviews (Table 3). Compared to persons interviewed by telephone, those interviewed in-person were younger, interviewed sooner after HIV diagnosis, and were more often male, non-white, men who have sex with men, and diagnosed at the PHSKC STD Clinic or an affiliated CBO ( $p < 0.001$ ). Compared to telephone interviews, in-person interviews resulted in significantly more partners per interviewed OP being named, notified, and newly diagnosed with HIV (Table 4). Adjusting for OP age, gender, race/ethnicity, sexual orientation, time from diagnosis to interview, type of diagnosing clinic, and DIS who performed the interview, in-person interviews were associated with a more than two-fold increase in HIV case-finding. There was no significant difference between in-person and telephone interviews in number of partners tested for HIV per OP interviewed.

Of the 358 in-person interviews, 27 were field investigations and 323 took place in the STD clinic. The remaining 8 in-person interviews did not have a location recorded.

We found fewer differences between field-based interviews and telephone interviews with no significant difference in the notification, HIV test, and HIV case-finding indices. Field-based interviews were associated with a lower contact index than telephone interviews (aRR 0.63, 95% CI 0.40-0.97). The evaluation of STD clinic interviews versus telephone interviews resulted in the same findings as our primary analysis. There were no associations found between OP age, gender, race/ethnicity, sexual orientation, and time from OP diagnosis to interview and the HIV case-finding index.

### *Sensitivity Analyses*

There were no substantial differences between verified and unverified outcomes. For in-person interviews, the verified syphilis brought-to-treatment index was 0.26 and the unverified brought-to-treatment index was 0.27. For telephone interviews, the verified syphilis brought-to-treatment index was 0.22 and the unverified brought-to-treatment index was 0.24. We found that all new HIV cases were verified, so the verified and unverified HIV case-finding indices were the same.

### **Discussion:**

In this retrospective cohort study of OPs who underwent PS interviews in King County, Washington from 2010-2014, we found that in-person PS interviews for syphilis were associated with identifying, notifying, testing, and empirically treating more partners compared to telephone PS interviews, but were not associated with identifying and treating more partners that actually had syphilis. In contrast, in-person HIV PS interviews were associated with better outcomes for almost all metrics, including HIV

case-finding. We found fewer differences between field-based in-person interviews and telephone interviews; however, the small number of field-based interviews conducted by our PS program likely limited the analysis as our study was not powered to investigate this comparison.

The overall case-finding indices and brought-to-treatment indices we report are consistent with those reported in prior studies. A study that compiled PS outcomes across the United States from 1976-2004 found that the median case-finding index for syphilis was 0.22 (range 0.05-0.46) and for HIV was 0.13 (range 0.03-0.75).<sup>12</sup> Both our overall case-finding indices - 0.27 and 0.07 for syphilis and HIV, respectively – and our separate case-finding indices for in-person and telephone interviews were within these ranges, suggesting that our program’s performance was similar to that reported in other areas during an earlier era.

We hypothesized that in-person interviews would be somewhat more effective than telephone interviews. However, focusing on the most important outcomes - case-finding and treatment of infected persons – in-person interviews were only superior for HIV. Some prior research suggests that in-person interviews are not always superior. Studies evaluating outcomes of in-person versus telephone interviews about sensitive topics such as alcohol abuse<sup>15</sup>, welfare use<sup>16</sup>, and food security<sup>17</sup>, found similar effectiveness of in-person and telephone interviews.

Onsite DIS and decreased time from diagnosis to interview have been associated with improved PS outcomes.<sup>18,19</sup> A study evaluating HIV PS outcomes in San Francisco found a case-finding index of 0.13 for OPs interviewed within 2 weeks of diagnosis compared to 0.05 for those conducted more than 2 weeks after diagnosis.<sup>20</sup>

Interestingly, our study found that in-person HIV PS interviews, which are typically performed by onsite DIS, occur a median of 10 days after diagnosis with a case-finding index of 0.10 and our telephone HIV PS interviews occur a median of 42 days after diagnosis with a case-finding index of 0.05. While time to interview may have influenced our results, the differences we observed in case-finding between in-person and telephone HIV PS interviews remained significant after adjusting for time to interview, suggesting that the timing of interviews did not entirely explain the differences we observed.

It is also possible that the significant psychosocial impact of receiving a new HIV diagnosis may be a reason why telephone interviews are not as effective as in-person interviews for HIV PS. A prior study found that HIV peer counselors perceive in-person interventions to be more effective than telephone interventions because face-to-face counseling allows for personal contact and facilitates trusting relationships.<sup>21</sup> Another study revealed that patients notified of their HIV status by telephone delayed establishing primary care considerably longer than those notified in person.<sup>22</sup>

The strengths of our study include the large, population-based sample size and the long-standing use of both in-person and telephone interviews for PS in our HIV/STD control program. However, our analysis has significant limitations. First and foremost, this is an observational study, and we cannot eliminate the possibility that our findings are influenced by residual confounding. Second, our results may have been influenced by ascertainment bias. Many persons diagnosed with STDs, including HIV, notify their partners without assistance from DIS. Indeed, we previously found that most partners identified through PS with newly diagnosed HIV infection were actually notified and

tested prior to being contacted by DIS.<sup>23</sup> It may be that in-person interviews do more to promote cases' willingness to name infected partners than true case identification. A randomized trial of different approaches to PS interviews would overcome the problem of residual confounding but might still be affected by ascertainment bias. To date, only one very small randomized controlled trial has evaluated HIV PS in the U.S., and we are not aware of any recent randomized trials related to syphilis PS.<sup>24</sup> Given the significant programmatic costs associated with PS and the many competing demands on DIS time, it may be worth conducting trials to better assess the true benefit of the more expensive traditional approach that focuses on in-person interviews. Lastly, our results may not be generalizable to other health jurisdictions. Predictors of PS success have been shown to vary by location.<sup>25</sup> Perhaps more importantly, very few of the in-person interviews undertaken by King County DIS involved field investigations; most persons were interviewed in our STD clinic, either because they were diagnosed there or came to the clinic after being diagnosed. As a result, our procedures were dissimilar from many areas where DIS initiate field investigations within days of learning of a new HIV or syphilis diagnosis.

Many U.S. health departments are trying to modernize their PS programs to evolve toward a broader model of field services. This effort requires reorganizing programs to confront a growing syphilis epidemic while simultaneously taking on new work providing HIV PS to all persons with newly diagnosed HIV infection, linking and relinking patients to HIV care, and promoting HIV pre-exposure prophylaxis. Success with this expanded portfolio of work will require willingness to reconsider traditional procedures. Our findings suggest that in-person interviews may not be superior to

telephone interviews for syphilis, at least in terms of syphilis case-finding and treatment of infected partners. While our results suggest that in-person interviews are better for HIV, insofar as they involve field investigations, they are almost certainly substantially more costly. If the resources required for in-person interviews lead to fewer persons with newly diagnosed HIV being interviewed, the cost of the intervention could offset its greater efficacy leading to lower public health impact. Ultimately, how to allocate scarce DIS resources requires defining the costs and outcomes associated with different bodies of work and then setting priorities. We believe that our findings support health departments adopting a more flexible approach to PS, one that allows for more routine use of telephone PS for syphilis, and at least some use of telephone PS as an initial means of contacting persons with newly diagnosed HIV infection. More rigorous, prospective studies of procedures are needed to define the most cost-effective methods for accomplishing the many tasks assigned to PS teams in the modern era of high impact HIV prevention.

## References:

1. O'Connor J, Matthews G. Informational privacy, public health, and state laws. *Am J Public Health*. 2011;101(10):1845-1850. doi: 10.2105/10.2105/AJPH.2011.300206 [doi].
2. Hogben M. Partner notification for sexually transmitted diseases. *Clin Infect Dis*. 2007;44 Suppl 3:S160-74. doi: CID40991 [pii].
3. Centers for Disease Control and Prevention (CDC). Recommendations for partner services programs for HIV infection, syphilis, gonorrhea, and chlamydial infection. *MMWR Recomm Rep*. 2008;57(RR-9):1-83; quiz CE1-4. doi: rr5709a1 [pii].
4. Hightow-Weidman L, Beagle S, Pike E, et al. "No one's at home and they won't pick up the phone": Using the internet and text messaging to enhance partner services in North Carolina. *Sex Transm Dis*. 2014;41(2):143-148. doi: 10.1097/OLQ.0000000000000087 [doi].
5. Kachur R, Adelson S, Firenze K, Herrera M. Reaching patients and their partners through mobile: Text messaging for case management and partner notification. *Sex Transm Dis*. 2011;38(2):149-150. doi: 10.1097/OLQ.0b013e3182036bab [doi].
6. Udeagu CC, Bocour A, Shah S, Ramos Y, Gutierrez R, Shepard CW. Bringing HIV partner services into the age of social media and mobile connectivity. *Sex Transm Dis*. 2014;41(10):631-636. doi: 10.1097/OLQ.0000000000000181 [doi].
7. Ehlman DC, Jackson M, Saenz G, et al. Evaluation of an innovative internet-based partner notification program for early syphilis case management, Washington, DC,

January 2007-June 2008. *Sex Transm Dis.* 2010;37(8):478-485. doi:  
10.1097/OLQ.0b013e3181e212cb [doi].

8. Mendez J, Maher J. Evidence supporting the use of text messaging for partner services. *Sex Transm Dis.* 2012;39(3):238-239. doi: 10.1097/OLQ.0b013e3182471b31 [doi].

9. Hogben M, Collins D, Hoots B, O'Connor K. Partner services in sexually transmitted disease prevention programs: A review. *Sex Transm Dis.* 2016;43(2 Suppl 1):S53-62. doi: 10.1097/OLQ.0000000000000328 [doi].

10. Hennessy M, Williams SP, Mercier MM, Malotte CK. Designing partner-notification programs to maximize client participation: A factorial survey approach. *Sex Transm Dis.* 2002;29(2):92-99.

11. Udeagu CC, Bocour A, Shepard CW. For partner services, do we need "face-time," or FaceTime?: Trends in relative proportion of in-person notifications and HIV testing after introduction of a telephone option for HIV exposure notification by public health. *Sex Transm Dis.* 2014;41(11):671-673. doi: 10.1097/OLQ.0000000000000192 [doi].

12. Brewer DD. Case-finding effectiveness of partner notification and cluster investigation for sexually transmitted diseases/HIV. *Sex Transm Dis.* 2005;32(2):78-83. doi: 00007435-200502000-00003 [pii].

13. King Holmes, MD, PhD, P. Sparling, MD, Walter Stamm, MD, Peter Piot, MD, PhD, Judith Wasserheit, MD, MPH, Lawrence Corey, MD, Myron Cohen, MD, D. Watts, MD.

Public health partner notification practices in high-income nations. In: *Sexually transmitted diseases*. 4th ed. McGraw-Hill; 2008.

14. StataCorp LP. Stata statistical software. 2015; Release 14.

15. Midanik LT, Greenfield TK. Telephone versus in-person interviews for alcohol use: Results of the 2000 national alcohol survey. *Drug Alcohol Depend*. 2003;72(3):209-214. doi: S0376871603002047 [pii].

16. Pridemore WA, Damphousse KR, Moore RK. Obtaining sensitive information from a wary population: A comparison of telephone and face-to-face surveys of welfare recipients in the United States. *Soc Sci Med*. 2005;61(5):976-984. doi: S0277-9536(05)00041-9 [pii].

17. Nord M, Hopwood H. Does interview mode matter for food security measurement? telephone versus in-person interviews in the current population survey food security supplement. *Public Health Nutr*. 2007;10(12):1474-1480. doi: S1368980007000857 [pii].

18. Taylor MM, Mickey T, Winscott M, James H, Kenney K, England B. Improving partner services by embedding disease intervention specialists in HIV-clinics. *Sex Transm Dis*. 2010;37(12):767-770. doi: 10.1097/OLQ.0b013e3181e65e8b [doi].

19. Rudy ET, Aynalem G, Cross J, Ramirez F, Bolan RK, Kerndt PR. Community-embedded disease intervention specialist program for syphilis partner notification in a clinic serving men who have sex with men. *Sex Transm Dis*. 2012;39(9):701-705. doi: 10.1097/OLQ.0b013e3182593b51 [doi].

20. Marcus JL, Bernstein KT, Klausner JD. Updated outcomes of partner notification for human immunodeficiency virus, San Francisco, 2004-2008. *AIDS*. 2009;23(8):1024-1026. doi: 10.1097/QAD.0b013e32832921a7 [doi].
21. Messias DK, Moneyham L, Murdaugh C, Phillips KD. HIV/AIDS peer counselors' perspectives on intervention delivery formats. *Clin Nurs Res*. 2006;15(3):177-196. doi: 15/3/177 [pii].
22. Samet JH, Freedberg KA, Stein MD, et al. Trillion virion delay: Time from testing positive for HIV to presentation for primary care. *Arch Intern Med*. 1998;158(7):734-740.
23. Golden MR, Stekler J, Kent JB, Hughes JP, Wood RW. An evaluation of HIV partner counseling and referral services using new disposition codes. *Sex Transm Dis*. 2009;36(2):95-101. doi: 10.1097/OLQ.0b013e31818d3ddb [doi].
24. Landis SE, Schoenbach VJ, Weber DJ, et al. Results of a randomized trial of partner notification in cases of HIV infection in North Carolina. *N Engl J Med*. 1992;326(2):101-106. doi: 10.1056/NEJM199201093260205 [doi].
25. Hoots BE, Lewis FM, Anschuetz G, et al. Would targeting increase efficiency of syphilis partner services programs?--data from New York City, Philadelphia, Texas, and Virginia. *Sex Transm Dis*. 2014;41(6):407-412. doi: 10.1097/OLQ.000000000000130 [doi].

Table 1. Characteristics of Original Patients with Early Syphilis Interviewed by Public Health - Seattle and King County Partner Services from 2010 to 2014, by Method of Interview (In-Person versus Telephone)

Original Patient Characteristic	Syphilis In-Person Interview N=682*	Syphilis Telephone Interview N=646	P-value
<u>Age</u>			
Median years (interquartile range)	37 (28-45)	41 (32-48)	<0.001
<u>Sex, n (%)</u>			
Male	668 (98)	627 (97)	0.299
Female	14 (2)	19 (3)	
<u>Race/ethnicity, n (%)</u>			
White	426 (63)	455 (70)	0.001
Black	79 (12)	48 (7)	
Latino	80 (12)	76 (12)	
Asian	25 (4)	15 (2)	
Other	49 (7)	26 (4)	
<u>Sexual orientation, n (%)</u>			
MSM	632 (93)	594 (92)	0.114
MSWE	36 (5)	27 (4)	
WSM	23 (2)	17 (3)	
Other	2 (<1)	8 (1)	
<u>HIV status, n (%)</u>			
Negative	358 (52)	226 (35)	<0.001
Positive	324 (48)	420 (65)	
<u>Location of interview, n (%)</u>			
STD clinic	624 (91)	-	
Field	26 (4)	-	
<u>Time to interview</u>			
Median days (interquartile range)	6 (0-14)	21 (12-34)	<0.001
<u>Place of diagnosis, n (%)</u>			
STD Clinic and/or MSM CBO	405 (59)	57 (9)	<0.001
MSM and/or HIV Specialty Clinic	125 (18)	304 (47)	
Non-Specialty Provider	152 (22)	285 (44)	
<u>Caseworker, n (%)</u>			
DIS #1	328 (48)	171 (26)	<0.001
DIS #2	91 (13)	275 (43)	
DIS #3	111 (16)	123 (19)	
DIS #4	6 (1)	44 (7)	
DIS #5	40 (6)	12 (2)	
Other DIS*	53 (8)	16 (2)	

HIV: Human immunodeficiency virus, STD: Sexually transmitted disease, MSM: Men who have sex with men, MSWE: Men who have sex with women exclusively, WSM: Women who have sex with men, CBO: Community based organization, DIS: Disease intervention specialist. \*Includes all other small volume caseworkers. Numbers may not sum to total because of missing data; all variables have  $\leq 5\%$  missing.

Table 2. Partner Services Indices for Early Syphilis in King County, WA 2010-2014

*Partner Services Indices	In-person Interview N=682 Counts Index	Telephone Interview N=646 Counts Index	**Unadjusted Relative Risk (95% CI)	P-value	***Adjusted Relative Risk (95% CI)	P-value
Contact index	2340 3.43	1329 2.06	1.67 (1.56, 1.78)	<0.001	1.68 (1.55, 1.82)	<0.001
Notification Index	1155 1.70	724 1.13	1.51 (1.38, 1.66)	<0.001	1.39 (1.24, 1.56)	<0.001
Syphilis test index	786 1.15	463 0.72	1.61 (1.43, 1.80)	<0.001	1.34 (1.16, 1.54)	<0.001
HIV test index	355 0.52	131 0.20	2.57 (2.10, 3.14)	<0.001	1.45 (1.14, 1.86)	0.003
Epidemiologic index	703 1.03	477 0.74	1.40 (1.24, 1.57)	<0.001	1.19 (1.03, 1.37)	0.017
Syphilis case-finding index	193 0.28	162 0.25	1.13 (0.92, 1.39)	0.257	0.91 (0.71, 1.17)	0.470
HIV case-finding index	14 0.02	5 0.01	2.65 (0.96, 7.36)	0.061	1.83 (0.56, 5.95)	0.316
Brought-to-treatment index	189 0.28	153 0.24	1.17 (0.95, 1.45)	0.149	0.93 (0.72, 1.21)	0.600

HIV: Human immunodeficiency virus

\*All indices have a denominator of total original patients interviewed (N). Numerators are:

Contact index: Partners named

Notification index: Partners notified

Syphilis test index: Partners tested for syphilis

HIV test index: Partners tested for HIV

Epidemiology index: Partners empirically treated for syphilis

Syphilis case-finding index: Partners diagnosed with syphilis

HIV case-finding index: Partners diagnosed with HIV

Brought-to-treatment index: Partners diagnosed with and treated for syphilis

\*\*Unadjusted Poisson regression model

\*\*\*Multivariable Poisson regression model adjusted for age, gender, race/ethnicity, sexual orientation, HIV status, time from diagnosis to interview, place of diagnosis, and disease intervention specialist who performed the interview.

Table 3. Characteristics of Original Patients with New HIV Diagnoses Interviewed by Public Health - Seattle and King County Partner Services from 2010 to 2014, by Method of Interview (In-Person versus Telephone)

Original Patient Characteristic	HIV In-Person Interview N=358	HIV Telephone Interview N=489	P-value
<u>Age</u>			
Median years (interquartile range)	32 (25-41)	36 (28-46)	<0.001
<u>Sex, n (%)</u>			
Male	333 (93)	430 (88)	0.015
Female	25 (7)	59 (12)	
<u>Race/ethnicity, n (%)</u>			
White	188 (53)	310 (64)	<0.001
Black	53 (15)	78 (16)	
Latino	63 (18)	61 (12)	
Asian	28 (8)	22 (5)	
Other	26 (7)	18 (4)	
<u>Sexual orientation, n (%)<sup>◊</sup></u>			
MSM	301 (84)	349 (72)	<0.001
MSWE	18 (5)	45 (9)	
WSM	20 (6)	50 (10)	
Other	6 (2)	8 (2)	
<u>Location of interview, n (%)</u>			
STD Clinic	323 (90)	-	
Field	27 (8)	-	
Other	8 (2)	-	
<u>Time to interview</u>			
Median days (interquartile range)	10 (5-18)	42 (24-72)	<0.001
<u>Place of diagnosis, n (%)</u>			
STD Clinic and/or MSM CBO	278 (78)	33 (7)	<0.001
MSM and/or HIV Specialty Clinic	7 (2)	78 (16)	
Non-Specialty Provider	73 (20)	378 (77)	
<u>Caseworker, n (%)</u>			
DIS #1	178 (50)	234 (48)	0.125
DIS #2	128 (36)	173 (35)	
DIS #3	18 (5)	40 (8)	
DIS #4	4 (1)	13 (3)	
DIS #5	2 (1)	6 (1)	
Other DIS*	28 (8)	23 (5)	

HIV: Human immunodeficiency virus, STD: Sexually transmitted disease, MSM: Men who have sex with men, MSWE: Men who have sex with women exclusively, WSM: Women who have sex with men, CBO: Community based organization, DIS: Disease intervention specialist. \*Includes all other small volume caseworkers. Numbers may not sum to total because of missing data. Variables have  $\leq 5\%$  missing unless indicated. <sup>◊</sup> 7.5% of telephone interviews had missing values for sexual orientation.

Table 4. Partner Services Indices for New HIV Diagnoses in King County, WA 2010-2014

*Partner Services Indices	In-person Interview N=358 Counts Index	Telephone Interview N=489 Counts Index	**Unadjusted Relative Risk (95% CI)	P-value	***Adjusted Relative Risk (95% CI)	P-value
Contact index	670 1.87	628 1.28	1.46 (1.31, 1.62)	<0.001	1.38 (1.18, 1.62)	<0.001
Notification Index	495 1.38	448 0.92	1.51 (1.33, 1.71)	<0.001	1.24 (1.03, 1.50)	0.026
HIV test index	219 0.61	235 0.48	1.27 (1.06, 1.53)	0.010	1.15 (0.88, 1.52)	0.303
HIV case-finding index	36 0.10	24 0.05	2.05 (1.22, 3.43)	0.006	2.17 (1.04, 4.50)	0.039

HIV: Human immunodeficiency virus

\*All indices have a denominator of total original patients interviewed (N). Numerators are:

Contact index: Partners named

Notification index: Partners notified

HIV test index: Partners tested for HIV

HIV case-finding index: Partners diagnosed with HIV

\*\*Unadjusted Poisson regression model

\*\*\*Multivariable Poisson regression model adjusted for age, gender, race/ethnicity, sexual orientation, time from diagnosis to interview, place of diagnosis, and disease intervention specialist who performed the interview.