

Primary Healthcare Staffing Needs Assessment—
A Discrete Event Simulation Study

Tolulope Abe

A thesis

submitted in partial fulfillment of the
requirement for the degree of

Master of Public Health

University of Washington

2014

Committee:

Kenneth Sherr

Richard Storch

Program Authorized to Offer Degree:

Health Services

©Copyright 2014

Tolulope Abe

University of Washington

Abstract

Primary Healthcare Staffing Needs Assessment—
A Discrete Event Simulation Study

Tolulope Abe

Chair of the Supervisory Committee:
Associate Professor Kenneth Sherr
Global Health

Background: Mozambique has a shortage of primary healthcare workers, effecting the quality of primary healthcare delivery. The specific aim of the proposed research is utilizing Industrial & Systems Engineering methods, specifically discrete event simulation, to identify strategies to improve primary healthcare delivery system performance in the Sofala Province of Mozambique. To improve primary healthcare delivery and assist facility-level management with decision-making, the research team decided to study staffing level effects on patient waiting time and develop a decision support tool that determines staffing level needs to maintain an average total wait time of less than 60minutes.

Methods: A discrete event simulation study was performed to model primary healthcare facility delivery systems using Arena Simulation software. What-if scenario experiments testing the impact health worker staffing level and patient demand fluctuations have on patient waiting time were designed using Statistical Analysis Software, and the experimental runs were performed using Arena Simulation Process Analyzer application. Minitab was used to perform a regression analysis to find mathematical models of wait time as a function of patient demand and staffing levels.

Results: The mathematical relationship of health worker staffing and patient demand levels with average patient wait time was estimated using regression analysis. The number of staff required to provide services was the aggregate of staffing needed for all patient types of a given facility. This approach produced staffing and wait time results that could not be validated and used to create a spreadsheet-based decision support tool.

Conclusion: The spreadsheet-based decision support tool aimed to bridge the gap between industrial and systems engineering methods and healthcare stakeholder knowledge of these methods while promoting implementation by allowing decision makers to perform simulations through a user-friendly tool. Although the proposed approach described in this study could not be validated, this method is beneficial when attempting to evaluate performance improvement strategy impacts on measures of performance. To create the spreadsheet-based decision support tool, it is recommended that an alternative approach be used to determine the relationship between wait time, patient demand, and staffing levels, such as queuing analysis.

TABLE OF CONTENTS

LIST OF FIGURES	i
LIST OF TABLES	ii
LIST OF ACRONYMS	iii
Introduction	1
Methods	4
Discussion	16
Conclusion	18
REFERENCES	19
Appendix A. Time Study Margin of Error Calculation	24
Appendix B. Conceptual Model: Process Flow Map of PHC Patient Flow	25
Appendix C. Conceptual Model: PHC Patient, Facility, & Worker Assumptions	26
Appendix D. DES Model Validation Hypothesis Test	27
Appendix E. DES Model Validation Hypothesis Test Results Summary	28
Appendix F. Rural Health Center Orthogonal Array for What-If Scenario Experiments	29
Appendix G. Urban Health Center Orthogonal Array for What-If Scenario Experiments	30
Appendix H. Rural Hospital Orthogonal Array for What-If Scenario Experiments	31
Appendix I. Spreadsheet-based Decision Support Tool Example	32
Appendix J. Regression Models and Statistical Summaries	34

LIST OF FIGURES

<i>Figure Number</i>		<i>Page</i>
Figure 1.	Simulation Study Framework	4
Figure 2.	Arena Process ANalyzer (PAN) Example	11
Figure 3.	Simulation Study Framework in the Global Health Setting	17

LIST OF TABLES

<i>Table Number</i>		<i>Page</i>
Table 1.	Data Collection Sample Size and Corresponding Margin of Error	6
Table 2.	Average Daily Patient Visits by Patient & Facility Type	7
Table 3.	Service Time Probability Distributions by Service & Facility Type	7
Table 4.	Healthcare Worker Volumes by Worker & Facility Type	8
Table 5.	DES Model Output Validated by System Expert	9
Table 6.	Current and What-if Scenario Staffing Levels by Worker and Facility Type	10

LIST OF ACRONYMS

ANC	Antenatal Care
DDCF	Doris Duke Charitable Foundation
DES	Discrete Event Simulation
GHI	Global Health Initiatives
HAI	Health Alliance International
ISE	Industrial and Systems Engineering
MOH	Ministry of Health
PAN	Process Analyzer
PHC	Primary Healthcare
RH	Rural Hospital
RHC	Rural Health Center
SAS	Statistical Analysis Software
UEM	University of Eduardo Mondlane
UHC	Urban Health Center
WHO	World Health Organization
WISN	Workload Indicators of Staffing Need

ACKNOWLEDGEMENTS

I would like to acknowledge the support of and express my gratitude to the following people who were instrumental to the completion of this thesis:

Kenneth Sherr, PhD

Thank you for your academic and career guidance. Your dedication to health systems research in the global health setting inspires me to continue research in the area.

Richard Storch, PhD

Thank you for your assistance throughout my graduate education. I appreciate the many research opportunities you have introduced to me—each of which deepened my understanding of Industrial & Systems Engineering applications in healthcare. Your mentorship has shaped my leadership and academic development.

Sarah Gimbel, PhD

Thank you for sharing your primary healthcare system knowledge during this research study. Your wealth of knowledge was invaluable to the completion of this study.

Operations Research Center in Beira, Mozambique

Thank you for serving as primary healthcare system experts and assisting with the data collection efforts. Without your tireless work in the field, this research study would not be possible.

My family

Your support has made my graduate education possible and worthwhile. Thank you for encouraging my academic and career pursuits.

DEDICATION

To Dad, Mom, James, & Michael

Introduction

In Mozambique, the mission to understand complex primary healthcare (PHC) delivery systems is a complicated task partially due to human and technical resource shortages. One critical issue with PHC delivery systems, in developing countries especially, is health worker shortages. Since 2000, Global Health Initiatives (GHIs) have been developed and implemented to control the spread of communicable diseases and strengthen health systems to improve overall population health [1]. Despite GHIs and their evidence-based strategies, health worker shortages are a barrier to program implementation, especially antiretroviral treatment services [1]. Africa has one of the most critical healthcare worker shortages, with the lowest number of healthcare workers per 10,000 people in 2013 [2]. Of the 57 countries worldwide experiencing health worker shortages in 2006, 36 were African countries [3]. In Mozambique specifically, from 2005-2012 there were 0.3 physicians and 3.4 nursing and midwifery personnel per 10,000 population compared to the average of 2.5 physicians and 9.1 nursing and midwifery personnel per 10,000 people in the African Region [2]. A resulting effect of these shortages is high disease burden.

In order to address health worker shortages, heuristic approaches have been employed to balance staffing levels with patient care demands. Previous approaches to staffing solutions include the World Health Organization's (WHO) Workload Indicators of Staffing Need (WISN) method [4] and modified versions of the WISN method [5]. WHO's WISN method uses current staffing levels of various healthcare workers, number of patients who utilize services provided by each type of healthcare worker, and average healthcare delivery service time data to determine staffing deficits and surpluses. WISN has been used in developing countries/regions, such as

Tanzania [6], India [7], Papua New Guinea [8], and East Africa [9]. Since facility, district, provincial, and national workload data may not be readily available, observational data collection, focus groups, and interviews with the assistance of doctors and nurses is required [5]. This can be a time-consuming process that interferes with accurate data collection while providing quality patient care.

In addition to WHO's WISN based calculations, Industrial & Systems Engineering (ISE) methods have been used to evaluate and manage staffing levels. ISE is defined as an engineering discipline:

...concerned with the design, improvement, and installation of integrated systems of men, materials, and equipment. It draws upon specialized knowledge and skill in the mathematical, physical, and social sciences, together with the principles and methods of engineering analysis and design, to specify, predict and evaluate the results to be obtained from such systems [10].

Interdisciplinary teams consisting of health care stakeholders, frontline workers, and industrial & systems engineers have collaborated to apply ISE methods to manufacturing and service industries. Per work published between 2000-2014 describing ISE applications to the hospital setting in the Americas and Europe, the most common ISE methods that have been used to evaluate and manage staffing levels are deterministic modeling [11-59] and discrete event simulation [19, 33, 46, 60-77]. Compared to WISN, simulation and optimization models may require a greater amount of detailed data to be collected in order to analyze health worker needs. As these systems grow increasingly complex, accurate modeling tools require a greater deal of

time and effort to create, validate, and verify [78]. Despite this, the aforementioned ISE methods remain powerful tools when analyzing complex health care systems.

Although ISE methods have been used in the healthcare industry of several industrialized countries, methods have been slower to spread in the healthcare industry of developing countries. This can be attributed to the lack of ISE awareness and knowledge. Another limitation of healthcare systems in both developed and developing countries is the ability to implement evidence-based performance improvement strategies. For simulation studies specifically, Brailsford suggests that overcoming this research challenge is based on the ability to balance a user-friendly simulation model with “scientific rigor and validity”. When describing the successful implementation of performance improvement strategies evaluated during a simulation study at a UK Hospital, he credits “the simplicity and interactive nature of the model” as a key factor for implementation [79].

The specific research question this study aims to answer is *how can discrete event simulation methods be used to determine primary healthcare staffing levels?* The objective of this study is to provide a description of a simulation study used to develop a decision support tool that determines healthcare staffing levels for PHC facilities in the Sofala Province in Mozambique. This study aims to provide 1) an industrial & systems engineering application in the primary health care setting and 2) a decision support tool to overcome user-friendliness barriers experienced when implementing improvement strategies observed during a simulation study. The spreadsheet-based decision support tool bridges the gap between ISE methods and PHC stakeholder and frontline staff knowledge of these methods.

Methods

The study method used to develop the PHC simulation model was based on a simulation study framework, illustrated in Figure 1 [80]. The steps in orange signify collaboration amongst simulation developers and PHC delivery system experts, while the steps in blue signify steps performed by simulation developers during this study. The chosen simulation method was discrete event simulation (DES). DES was chosen due to its advantages, which “stems from its flexibility, as well as the ability to handle variability, uncertainty, and complexity. It allows for an assessment of efficiency of a given system, performance of the ‘what-if’ analysis, as well as




Figure 1. Simulation Study Framework

the possibility to design a new system” [81].

The following method subsections based on the simulation study framework steps describe the simulation study and spreadsheet-based decision support tool development.

Formulate Problem & Plan Study

To formulate the problem, the research team first set the objectives of the study: 1) strengthen integrated health systems management; 2) develop appropriate tools to facilitate decision-making; and 3) build capacity for conducting innovative ISE studies and operations research to guide system-strengthening efforts. In order to meet these objectives, the project team proposed the development of a DES to study the relationship between patient demand, health

worker volume, and average total patient wait time in the PHC setting. Average total patient

wait time, the main performance measure of the study, is defined as the total amount of time a patient waits for services during a single PHC visit—from the time a patient arrives to a facility to the time that the patient leaves the facility—averaged for all patients of a given type.

Due to its poor health outcomes and high HIV prevalence, Sofala province was selected as the setting for this study. Sofala province has a variety of healthcare facilities offering different healthcare services. A simplifying assumption was that the majority of PHC services are sought from rural hospitals (RH), rural health centers (RHC), and urban health centers (UHC). The types of patients who typically seek PHC services at these facilities include adults receiving outpatient consultations, children receiving immunizations and/or outpatient consultations, and pregnant women receiving their first antenatal care. Additionally, laboratory and pharmacy services are available at each of the three facility types.

Define Model & Collect Data

A time study and health worker interviews were performed to collect service time data and define the conceptual model. The target population during the study were all patients who visit a rural health center, urban health center or rural hospital in the Sofala Province of Mozambique for PHC services. It was assumed that these patients were commonly adult and pediatric outpatients, some of whom require immunizations (well child patients), and pregnant patients seeking their first antenatal care visit (1st ANC patients). Data collection was performed on selected days during January – April 2011. To perform the on-site time study, data collectors were stationed at each service area and documented the start and end times for each service provided by health workers and times between services. Due to the limited number of data

collectors, data was collected on two-four separate days, typically during the morning. Based on the data collected, the following is the margin of error for each patient type at each facility where data was collected (Appendix A). Table 1 shows the sample size and margin of error (95% confidence level) for each service at each health facility.

Table 1. Data Collection Sample Size and Corresponding Margin of Error

	Outpatient			Well Child			Antenatal			Laboratory			Pharmacy		
	<i>n</i>	<i>st.dev.</i>	<i>E</i>	<i>n</i>	<i>st.dev.</i>	<i>E</i>	<i>n</i>	<i>st.dev.</i>	<i>E</i>	<i>n</i>	<i>st.dev.</i>	<i>E</i>	<i>n</i>	<i>st.dev.</i>	<i>E</i>
Rural Health Center	63	30.70	7.58	34	5.20	1.75	27	13.49	5.09	90	6.06	1.25	176	9.39	1.39
Urban Health Center	50	27.81	7.71	79	40.68	8.97	61	28.71	7.20	219	14.10	1.87	162	9.87	1.52
Rural Hospital	54	20.17	5.38	56	46.75	12.24	56	40.47	10.60	197	17.54	2.45	286	27.21	3.15

The sample size (*n*) for each service ranged from 27 to 286 observations, while the margin of error (*E*) ranged from 1.25 to 12.24 (assuming the sample is normally distributed). This may mean that the sample collected for outpatient, well child, and antenatal services did not accurately represent the population of patients who seek these services.

A conceptual model in the form of a basic process flow map (Appendix B) and list of assumptions (Appendix C) was created using the information gathered during the health worker interviews and time study. The basic process flow map illustrates the services each patient type received during a typical visit at each health facility. The list of assumptions provides more detail regarding patient flow, facility operations, and health worker staffing.

Data Validation

To validate that the data collected was an accurate representation of the service and waiting time experienced by patients, descriptive statistics of the collected data were presented to PHC frontline staff, managers, and stakeholders. Those who reviewed the data confirmed its accuracy.

Construct Computer Simulation Model

Model inputs and patient flow logic are required to construct the DES model. The model inputs for this simulation study were average number of arrivals that occur for each patient type during a workday, the service time probability distributions for each service, the patient flow logic, and the number and type of healthcare workers who provide PHC services. The software used to construct the DES model was Rockwell Automation’s Arena® Simulation Software (v13.2).

A 2012 census report was the data source used to estimate the average daily visits for each patient type at each facility. Table 2 indicates the average daily visits used as an input for the model.

Table 2. Average Daily Patient Visits by Patient & Facility Type

	Average Daily Visits		
	Outpatients ‡	Immunization Patients	1st Antenatal Care Patients
Rural Health Center	75	17	4
Rural Hospital	351	127	14
Urban Health Center	912	102	19

‡Per assumption, 70% of outpatients are adults, 30% are pediatric

The data collected during the time study was used to determine the service time probability distributions for each service. Arena’s® Input Analyzer application was used to find a probability distribution that best fit the observed data. Table 3 displays the service time distributions used in the model, measured in minutes unless otherwise stated.

Table 3. Service Time Probability Distributions by Service & Facility Type

	Service Time Probability Distributions		
	Rural Health Center	Rural Hospital	Urban Health Center
Registration	n/a	Uniform(0,15)†	Uniform(0,15)†
Adult Outpatient Consultation	0.5 + LOGN(3.51, 2.81)	0.5 + LOGN(7.26, 6.15)	TRIA(0.5, 2.12, 9.5)
Pediatric Outpatient & Well Child Outpatient Consultation	0.5 + LOGN(3.51, 2.81)	0.5 + LOGN(7.26, 6.15)	TRIA(0.5, 2.12, 9.5)
Immunizations	0.5 + LOGN(1.57, 1.58)	0.5 + LOGN(4.25, 4.37)	0.5 + 7 * BETA(0.653, 3.53)
Laboratory Services	0.5 + 35 * BETA(0.525, 2.86)	0.5 + 56 * BETA(0.649, 2.59)	0.5 + 53 * BETA(0.71, 1.8)
Pharmacy Services Drop Off	Uniform(0.5, 1)	Uniform(0.5, 1)	Uniform(0.5, 1)
Pharmacy Service Delivery	Uniform(0.5, 1.5)	Uniform(0.5, 1.5)	Uniform(0.5, 1.5)
First Antenatal Care Visit	1.5 + 19 * BETA(0.831, 1.52)	POIS(8.85)	1.5 + LOGN(6.6, 8.3)

† Measured in seconds

The patient flow logic of the model was based on the conceptual model process flow and PHC delivery system assumptions (Appendix B & C). Table 4 displays the number of healthcare workers at each facility based on a report of healthcare workers allocated to each healthcare facility.

Table 4. Healthcare Worker Volumes by Worker & Facility Type

Healthcare Worker Types and Volumes				
	<i>Services Provided</i>	<i>Rural Health Center</i>	<i>Rural Hospital</i>	<i>Urban Health Center</i>
Registration Agent	<i>give patient a service number as they arrive</i>	n/a	3	1
Outpatient Nurse	<i>provide outpatient treatment</i>	2	14	9
Outpatient Doctor	<i>provide outpatient treatment</i>	n/a	1	1
Immunization Agent	<i>immunize well child care patients and float</i>	1	2	2
Antenatal Care Nurse	<i>provide 1st antenatal care services</i>	2	5	5
Laboratory Worker	<i>provide laboratory services</i>	1	2	4
Pharmacy Worker	<i>provide pharmacy services</i>	1	3	6

Perform a Pilot Run, Verification and Validation

Once construction of the model was complete, the model was run so that output data could be verified and validated. While constructing the model, patients were “animated”, or in other words assigned a picture, so patient flow through a PHC facility could be observed while the model was running to verify the patient logic was correctly constructed in the simulation model. To validate that the simulation model accurately represents the actual PHC delivery services at a given facility, hypothesis tests were performed comparing simulation output data results to actual system results. The performance measure compared for both the actual system and the model was the total time each patient type spends waiting during a visit, which is the performance measure of interest. This measure was not originally collected during the time study, so additional data was collected at each facility where the original data was collected and during the same months in order to minimize variability. Hypothesis testing comparing actual to model average total patient wait time found that the actual and model wait times are statistically different for each patient type at each facility except for 1st ANC patients who received care in a

Urban Health Center (Appendix D & E). This may be a result of the small sample size of the time study data collected due to limited data collection resources. A system expert reviewed and approved the conceptual model and summary wait times and concluded that although the model and time study data are statistically different, they had observed patient wait times in the field similar to the simulation model output. Table 5 summarizes the model output, including average total wait times, approved by the system expert.

Table 5. DES Model Output Validated by System Expert

	% Requiring Lab and/or Pharmacy	Avg. Daily Arrivals	Avg Daily Departures	% LWBS	Total Wait Time (min)	Total Visit Time (min)
RHC Adult Outpatient	60	53	52	2%	184	193
RHC Pediatric Outpatient	45	23	22	4%	115	123
RHC 1st ANC Patient	10	4	4	0%	42	51
RHC Well Child Patient	0	17	17	0%	47	50
UHC Adult Outpatient	60	639	509	20%	197	205
UHC Pediatric Outpatient	45	274	186	32%	183	190
UHC 1st ANC Patient	10	19	19	0%	44	53
UHC Well Child Patient	0	102	101	1%	242	246
RH Adult Outpatient	60	246	221	10%	164	177
RH Pediatric Outpatient	45	106	95	10%	143	155
RH 1st ANC Patient	10	14	14	0%	42	51
RH Well Child Patient	0	127	127	0%	160	167

Design and Perform Experiments

Following the validation, the simulation model was used to study what-if scenarios. In order to create the decision support tool that would provide information on the number of healthcare workers required as average daily visits fluctuated while aiming to reduce average total wait time, a simulation modeling experiment was designed. The what-if scenario tests determine average total wait time resulting from varying healthcare worker levels and patient demand. For the varying patient demand, average daily visits were increased by up to 100% and decreased by down to 90% at 10% intervals, making a total of 20 patient demand levels tested, including the current patient demand level (0% increase/decrease). The staffing level range for each type of healthcare worker was set at 6, for this was determined to be a potentially feasible staffing level

range that could be fulfilled if necessary. Table 6 features the current and possible staffing minimum and maximum number of healthcare workers for each healthcare worker and facility type.

Table 6. Current and What-if Scenario Staffing Levels by Worker and Facility Type

	Rural Health Center Experiment Staffing Levels			Urban Health Center Experiment Staffing Levels			Rural Hospital Experiment Staffing Levels		
	Current	Min	Max	Current	Min	Max	Current	Min	Max
Registration Agent (Primary)	n/a	n/a	n/a	1	1	6	2	1	6
Registration Agent (Secondary)	n/a	n/a	n/a	n/a	n/a	n/a	1	1	6
Outpatient RN	2	1	6	9	7	12	14	12	17
Outpatient MD	n/a	n/a	n/a	1	1	6	1	1	6
ANC (ESMI) RN	2	1	6	5	3	8	5	3	8
Pharm Worker	1	1	6	6	4	9	3	1	6
Lab Worker	1	1	6	4	5	7	2	1	6
Immunization (Pre) Tech	1	1	6	2	1	6	2	1	6
	5 factors, 6 levels = 6⁵ = 7,776 Experiments			7 factors, 6 levels = 6⁷ = 279,936			8 factors, 6 levels = 6⁸ = 1,679,616		

A single experiment is a combination of different levels of each healthcare worker to assess their impact on average total wait time at a specific patient demand level. A full-randomized experiment testing every combination of the staffing levels would require a total of 1,967,328 randomized experiments for each patient demand level at all three facilities. To reduce the number of experiments, balanced orthogonal arrays that would produce the similar statistical conclusions with fewer experiments were found using SAS Statistical Analysis Software (v9.4) (Appendix F-H). A total of 255 experiments for each volume level at all three facilities were designed using SAS.

Analyze Output Data

Arena’s Process Analyzer (PAN) application was used to perform the what-if scenario experiments. Figure 2 displays an example of the first 25 experiments run in PAN for a Rural Health Center at the current (0% increase/decrease) patient demand level.

Scenario Properties			Controls					Response Variables				
S	Name	Program File	Reps	Nurse of CS Chibabava	ESMI of CS Chibabava	PharAgent of CS Chibabava	LabAgent of CS Chibabava	PreTech of CS Chibabava	Adult_Patient of CS Chibabava WaitTime	Pediatric Outpatient of CS Chibabava Wait Time	Pregnant_Patient of CS Chibabava Wait Time	Well Child Patient of CS Chibabava Wait Time
1	Current Patient Volume	2: Moz Healthcare Delivery System Simulation Model.6.17.14.p	30	2	2	1	1	1	136	96	38	46
2	Current Patient Volume	2: Moz Healthcare Delivery System Simulation Model.6.17.14.p	30	1	1	4	4	2	215	141	45	41
3	Current Patient Volume	2: Moz Healthcare Delivery System Simulation Model.6.17.14.p	30	3	6	5	6	4	100	74	33	35
4	Current Patient Volume	2: Moz Healthcare Delivery System Simulation Model.6.17.14.p	30	4	5	4	1	3	91	70	33	36
5	Current Patient Volume	2: Moz Healthcare Delivery System Simulation Model.6.17.14.p	30	2	4	2	6	2	129	91	33	39
6	Current Patient Volume	2: Moz Healthcare Delivery System Simulation Model.6.17.14.p	30	1	6	1	2	1	214	141	32	48
7	Current Patient Volume	2: Moz Healthcare Delivery System Simulation Model.6.17.14.p	30	5	1	1	1	3	86	68	48	35
8	Current Patient Volume	2: Moz Healthcare Delivery System Simulation Model.6.17.14.p	30	2	5	1	5	5	131	92	31	35
9	Current Patient Volume	2: Moz Healthcare Delivery System Simulation Model.6.17.14.p	30	2	5	2	4	6	132	93	33	35
10	Current Patient Volume	2: Moz Healthcare Delivery System Simulation Model.6.17.14.p	30	5	3	5	1	6	82	67	34	33
11	Current Patient Volume	2: Moz Healthcare Delivery System Simulation Model.6.17.14.p	30	2	2	6	4	2	129	90	35	39
12	Current Patient Volume	2: Moz Healthcare Delivery System Simulation Model.6.17.14.p	30	3	3	6	4	3	102	75	34	36
13	Current Patient Volume	2: Moz Healthcare Delivery System Simulation Model.6.17.14.p	30	3	1	5	5	1	101	75	45	45
14	Current Patient Volume	2: Moz Healthcare Delivery System Simulation Model.6.17.14.p	30	3	2	4	5	6	102	74	36	34
15	Current Patient Volume	2: Moz Healthcare Delivery System Simulation Model.6.17.14.p	30	6	5	3	1	1	79	66	33	44
16	Current Patient Volume	2: Moz Healthcare Delivery System Simulation Model.6.17.14.p	30	4	5	6	5	2	84	67	32	38
17	Current Patient Volume	2: Moz Healthcare Delivery System Simulation Model.6.17.14.p	30	1	6	6	3	6	213	140	33	37
18	Current Patient Volume	2: Moz Healthcare Delivery System Simulation Model.6.17.14.p	30	2	1	3	6	3	129	90	45	37
19	Current Patient Volume	2: Moz Healthcare Delivery System Simulation Model.6.17.14.p	30	4	2	1	4	5	89	71	35	34
20	Current Patient Volume	2: Moz Healthcare Delivery System Simulation Model.6.17.14.p	30	6	5	4	2	6	74	61	32	33
21	Current Patient Volume	2: Moz Healthcare Delivery System Simulation Model.6.17.14.p	30	5	2	4	6	4	77	64	38	33
22	Current Patient Volume	2: Moz Healthcare Delivery System Simulation Model.6.17.14.p	30	4	3	6	6	6	86	67	35	33
23	Current Patient Volume	2: Moz Healthcare Delivery System Simulation Model.6.17.14.p	30	5	5	5	3	2	77	64	33	37
24	Current Patient Volume	2: Moz Healthcare Delivery System Simulation Model.6.17.14.p	30	2	2	5	5	1	134	95	35	46
25	Current Patient Volume	2: Moz Healthcare Delivery System Simulation Model.6.17.14.p	30	1	5	2	6	3	214	142	32	39

Figure 2. Arena Process ANalyzer (PAN) Example

As shown in figure 2, the controls of the experiment are the combinations of healthcare worker staffing levels in the middle of the figure and the resulting average total wait time for each patient type are the experiment response variables at the right end of the figure. The what-if scenario experiments produced response data that can be used to determine the relationship between average total wait time, healthcare worker levels, and patient demand.

Document, Present, and Implement Results

The next step of the study was to create a decision support tool using the resulting what-if scenario experiment data. Once the relationship between average total patient wait time, health worker staffing levels, and patient demand was estimated, the staffing levels could be determined based on predicted average daily visits and a targeted average total patient wait time less than 60min. The aim of developing the decision support tool was to create a user-friendly interface accessible via any computer with Microsoft Excel macros-running capability.

The tool prompts a PHC staffing decision maker to select their given PHC health facility type, specify the expected number of visits for each patient type for an upcoming workday and a targeted average total wait time for all patient types (60min), and the macros outputs the number of healthcare workers based on the staffing level statistical relationship with these criteria (Appendix I).

To determine the statistical relationship between average total wait time, healthcare worker staffing level, and patient demand, Minitab Statistical Software was used to perform regression analysis and determine the regression models that would be used to create the decision support tool. A multiple regression analysis was performed for each patient type at each PHC facility. For the regression analysis performed for each patient type, patient demand and all healthcare workers who provide PHC services were the regressor variables for the regression model, and their average total wait times were the response variables. If the results showed that the p-values of the hypothesis test were greater than the selected $\alpha=0.05$, these independent variables were removed and the regression analysis was performed again, a stepwise regression analysis process known as backward elimination. Additionally, in order to meet the assumptions of regression analysis—residuals fit a $N(0,\sigma^2)$ distribution and residuals are independent of each other—a Box-Cox optimal power transformation was performed. Backward elimination of transformed regression models was performed when necessary as previously described. Based on backward elimination and transforming the response variable when needed, a total of 41 regression models were created. Minitab provided statistics to test the adequacy of each model. To determine the models that would be used for the decision support tool, the following statistics were analyzed to assess each regression models adequacy:

- ANOVA F-Test: Test how well independent variables predicts the response variable;
target: p-value < 0.05
- Lack-of-Fit: test how well the model fits the data; *target: p-value > 0.05*
- R² adjusted: percent of response variable variation explained by the independent variable;
target: larger the better
- Durbin Watson Statistic: test of residual correlation; *target: approximately equal to 2*
- Anderson Darling Statistic & p-value- hypothesis test of how well the residuals fit a normal distribution; *target: Anderson-Darlin Statistic for the smaller the better; p-value > 0.05*

The statistics for each model were summarized and analyzed to determine the best regression model to use for the decision support tool [Appendix J]. None of the regression models met all target criteria, including models with an optimal Box-Cox transformation of the response variable. Therefore, regression models to include in the decision support tool were chosen for each patient type if they met most of the model adequacy target requirements listed above. Based on these criteria, the regression models most likely to meet all statistical requirements were regression models where the response variables were transformed via optimal Box-Cox transformation including all healthcare workers who provide services to the given patient type as the regressors. The exception to this was adult outpatient average total wait time at Urban Health Centers where the regression model chosen was the transformed model with backward elimination and well child care patients at Rural Hospitals where the regression model chosen was a non-transformed model with backward elimination.

To calculate the number of healthcare workers based on patient demand and average total wait time targets, the selected regression models were used as functions in a Microsoft Excel spreadsheet. The different staffing levels tested during the what-if scenario analysis served as the values for the independent variables of the regression models in the spreadsheet. Once average daily visit data is entered into the decision support tool for each patient type, the staffing level for each patient type was determined by finding the staffing level scenario resulting in the smallest difference between the expected average total wait time and the calculated average total wait time. For the selected health facility, the number of health workers needed is the aggregate of staffing levels for all patient types. The decision support tool outputs the aggregate staffing levels.

The average daily visits, healthcare worker volumes, and resulting average total wait time data from twenty randomly selected what-if scenarios were used to test the validity of the decision support tool for each health facility. The patient volumes and wait times for each individual patient type at each individual PHC facility were entered directly into the regression models. The staffing levels output by the decision support tool was compared to the what-if scenarios. The regression models could not be validated, meaning that staffing levels based on wait times and patient demand levels calculated with the regression models was not similar to the staffing levels for the wait times and patient demand levels found during the what-if scenario.

Upon further review of this method, the researchers have decided that identifying staffing needs for each facility based on average service time and average daily visits while controlling average total patient wait time can be performed using queuing analysis instead of a DES study. DES

was best used in this study to identify services where patients experience higher wait times and test strategies to alleviate these bottlenecks, like block scheduling strategies to smooth arrival patterns throughout the workday or reallocating a nurse to assist with registration during the peak hours of the day.

Study Limitations

The regression models used in the decision support tool are based on DES output data that was validated based on system expert approval, but the output data was not statistically valid. This may be due to the small sample size of data collected during the time study. Although the smaller sample size was attributed to the shortages in data collection resources during this study, future researchers interested in simulation studies in the healthcare setting should consider collecting a larger sample size. Additionally, a small sample of data should be set aside so it is not used as an input into the DES model and can therefore be used to validate the DES model output.

Another limitation of this study is that if any PHC system changes occur beyond changes in staffing levels and average daily volumes, such as PHC delivery performance improvement changes that impact service times or assumptions, the DES model would need to be updated to capture how these system changes impact average total patient wait times. In this case, a DES developer or healthcare decision maker familiar with DES and regression analysis is required to make decision support tool changes.

Discussion

This study presented a simulation study performed remotely where the results were used to develop a spreadsheet-based decision support tool. The decision support tool aimed to identify the staffing level required to provide PHC services to an expected number of daily patient visits while maintaining a targeted average total patient wait time. The next step in this study is to use a different method to calculate staffing level needs for the decision support tool and implement the decision support tool amongst PHC facility management in the Sofala province.

To successfully implement a decision support tool in the PHC setting using the originally proposed method described in this study, the results of the simulation and the output featured in the study need to be reviewed with multiple PHC stakeholders, including frontline staff, at the provincial level to garner feedback to improve the decision support tool as needed as well as facilitate wide-spread support and utilization.

A limitation of the study is that frontline staff and end users were not involved in each step of the simulation study. Their involvement throughout the course of the study could assist with validation of the model as well as facilitate successful implementation of the decision support tool. To overcome this limitation for future studies, figure 3 proposes a simulation study in the healthcare setting framework based on literature of suggestions for successful simulation study implementation [80, 82].




Figure 3. Simulation Study Framework in the Global Health Setting

The revised framework proposes that a steering committee made up of end users, decision makers, and problem owners are involved with each step of the simulation study. During this study, periodic meetings for data and model validation and for study status updates were held with system experts off-site, however future studies may consider performing these meetings on-site to promote study engagement amongst on-site steering committee members, which will ultimately increase implementation successfulness.

Conclusion

This study aimed to develop a spreadsheet-based decision support tool accessible through Microsoft Excel. The tool would determine staffing needs based on the statistical relationship between staffing levels, patient demand, and wait time. Data produced from a discrete event simulation study was used to find the relationship amongst these variables. With a user-friendly interface, this decision support tool would support the implementation of industrial and systems engineering approaches in the primary health care setting.

The method proposed in this study could not be validated. The researcher suggests that future efforts to develop a decision support tool that determines staffing needs based on wait time and patient demand should use queuing analysis methods. To facilitate implementation of a revised decision support tool, end users, decision makers, and problem owners should be involved with each stage of the study and decision support tool development. Future work should determine how the decision support tool can be made more flexible to accurately capture structural changes, staffing level changes, and average daily visit fluctuations beyond the values tested during this study. Additionally, future work comparing the results from this study with other staffing methods, such as WISN and deterministic modeling, should be performed and reviewed with stakeholders to assess which method produces feasible staffing levels while achieving performance targets.

REFERENCES

1. Biesma, R.G., et al., *The effects of global health initiatives on country health systems: a review of the evidence from HIV/AIDS control*. Health policy and planning, 2009. **24**(4): p. 239-252.
2. WHO, *World health statistics 2013.*, in *WHO Library Cataloguing-in-Publication Data*. 2013, World Health Organization: Geneva, Switzerland.
3. Callaghan, M., N. Ford, and H. Schneider, *Review A systematic review of task-shifting for HIV treatment and care in Africa*. Human resources for health, 2010. **8**: p. 8-16.
4. Shipp, P.J., *Workload indicators of staffing need (WISN): a manual for implementation*. 1998: WHO Division of Human Resources Development and Capacity Building Geneva, Switzerland.
5. Hontelez, J., et al., *Human resources needs for universal access to antiretroviral therapy in South Africa: a time and motion study*. Hum Resour Health, 2012. **10**(1): p. 39.
6. Nyamtema, A., et al., *Staffing needs for quality perinatal care in Tanzania*. African journal of reproductive health, 2008. **12**(3).
7. Hagopian, A., et al., *Applying WHO's 'workforce indicators of staffing need'(WISN) method to calculate the health worker requirements for India's maternal and child health service guarantees in Orissa State*. Health policy and planning, 2011: p. czr007.
8. Kolehmainen-Aitken, R.-L. and P. Shipp, *'Indicators of staffing need': assessing health staffing and equity in Papua New Guinea*. Health Policy and Planning, 1990. **5**(2): p. 167-176.
9. Musau, P., et al., *Workload Indicators of Staffing Need method in determining optimal staffing levels at Moi Teaching and Referral Hospital*. East African medical journal, 2008. **85**(5): p. 232-239.
10. Hodson, W.K., *Maynard's industrial engineering handbook*. 4 ed. 1992, New York City, New York: McGraw-Hill.
11. Aickelin, U. and K.A. Dowsland, *Exploiting problem structure in a genetic algorithm approach to a nurse rostering problem*. Journal of Scheduling, 2000. **3**: p. 139-153.
12. Beaulieu, H., et al., *A mathematical programming approach for scheduling physicians in the emergency room*. Health Care Manag Sci, 2000. **3**(3): p. 193-200.
13. Dowsland, K.A. and J.M. Thompson, *Solving a Nurse Scheduling Problem with Knapsacks, Networks and Tabu Search*. The Journal of the Operational Research Society, 2000. **51**(7): p. 825-833.
14. Gascon, V., et al., *Scheduling the flying squad nurses of a hospital using a multi-objective programming model*. Annals of Operations Research, 2000. **96**: p. 149-166.
15. Valouxis, C. and E. Housos, *Hybrid optimization techniques for the workshift and rest assignment of nursing personnel*. Artif Intell Med, 2000. **20**(2): p. 155-75.
16. Athanassopoulos, A. and C. Gounaris, *Assessing the technical and allocative efficiency of hospital operations in Greece and its resource allocation implications*. European Journal of Operational Research, 2001. **133**(2): p. 416-431.
17. Ferland, J., et al., *Generalized Assignment Type Goal Programming Problem: Application to Nurse Scheduling*. Journal of Heuristics, 2001. **7**(4): p. 391-413.

18. Blake, J.T., F. Dexter, and J. Donald, *Operating Room Managers' Use of Integer Programming for Assigning Block Time to Surgical Groups: A Case Study*. *Anesth Analg*, 2002. **94**: p. 143-148.
19. Chan, S.Y., et al., *Operations research methods applied to workflow in a medical records department*. *Health Care Manag Sci*, 2002. **5**(3): p. 191-9.
20. Mullinax, C. and M. Lawley, *Assigning patients to nurses in neonatal intensive care*. *Journal of the Operational Research Society*, 2002. **53**(1): p. 25-35.
21. Sherali, H.D., M.H. Ramahi, and Q.J. Saifee, *Hospital resident scheduling problem*, *Production Planning & Control*. *The Management of Operations*, 2002. **13**(2): p. 220-233.
22. Cheang, B., et al., *Nurse rostering problems--a bibliographic survey*. *European Journal of Operational Research*, 2003. **151**(3): p. 447-460.
23. Aickelin, U. and K.A. Dowsland, *An indirect Genetic Algorithm for a nurse-scheduling problem*. *Computers & Operations Research*, 2004. **31**(5): p. 761-778.
24. Blöchliger, I., *Modeling staff scheduling problems. A tutorial*. *European Journal of Operational Research*, 2004. **158**(3): p. 533-542.
25. Bard, J.F. and H.W. Purnomo, *A column generation-based approach to solve the preference scheduling problem for nurses with downgrading*. *Socio-Economic Planning Sciences*, 2005. **39**(3): p. 193-213.
26. Bard, J.F. and H.W. Purnomo, *Preference scheduling for nurses using column generation*. *European Journal of Operational Research*, 2005. **164**(2): p. 510-534.
27. Day, T.E., J.T. Napoli, and P.C. Kuo, *Scheduling the resident 80-hour work week: an operations research algorithm*. *Curr Surg*, 2006. **63**(2): p. 136-41; discussion 141-2.
28. Topaloglu, S., *A multi-objective programming model for scheduling emergency medicine residents*. *Computers & Industrial Engineering*, 2006. **51**(3): p. 375-388.
29. Bard, J.F. and H.W. Purnomo, *Cyclic preference scheduling of nurses using a Lagrangian-based heuristic*. *Journal of Scheduling*, 2007. **10**(1): p. 5-23.
30. Beliën, J. and E. Demeulemeester, *Building cyclic master surgery schedules with leveled resulting bed occupancy*. *European Journal of Operational Research*, 2007. **176**(2): p. 1185-1204.
31. Bester, M.J., I. Nieuwoudt, and J.H. Van Vuuren, *Finding good nurse duty schedules: a case study*. *Journal of Scheduling*, 2007. **10**(6): p. 387-405.
32. Gutjahr, W.J. and M.S. Rauner, *An ACO algorithm for a dynamic regional nurse-scheduling problem in Austria*. *Computers & Operations Research*, 2007. **34**(3): p. 642-666.
33. Odegaard, F., et al., *Improving the efficiency of hospital porter services*. *J Healthc Qual*, 2007. **29**(1): p. 12-18.
34. Parr, D. and J.M. Thompson, *Solving the multi-objective nurse scheduling problem with a weighted cost function*. *Annals of Operations Research*, 2007. **155**(1): p. 279-288.
35. Sinreich, D. and O. Jabali, *Staggered work shifts: a way to downsize and restructure an emergency department workforce yet maintain current operational performance*. *Health Care Management Science*, 2007. **10**(3): p. 293-308.
36. Beliën, J. and E. Demeulemeester, *A branch-and-price approach for integrating nurse and surgery scheduling*. *European Journal of Operational Research*, 2008. **189**(3): p. 652-668.

37. Ogulata, S.N., M. Koyuncu, and E. Karaskas, *Personnel and patient scheduling in the high demanded hospital services: a case study in the physiotherapy service*. J Med Syst, 2008. **32**(3): p. 221-8.
38. Aickelin, U., E.K. Burke, and J.P. Li, *An Evolutionary Squeaky Wheel Optimization Approach to Personnel Scheduling*. Ieee Transactions on Evolutionary Computation, 2009. **13**(2): p. 433-443.
39. Cohn, A., et al., *Scheduling Medical Residents at Boston University School of Medicine*. Interfaces, 2009. **39**(3): p. 186-195.
40. De Grano, M.L., D.J. Medeiros, and D. Eitel, *Accommodating individual preferences in nurse scheduling via auctions and optimization*. Health Care Manag Sci, 2009. **12**(3): p. 228-42.
41. Burke, E.K., J. Li, and R. Qu, *A hybrid model of integer programming and variable neighbourhood search for highly-constrained nurse rostering problems*. European Journal of Operational Research, 2010. **203**(2): p. 484-493.
42. Roland, B., et al., *Scheduling an operating theatre under human resource constraints*. Computers & Industrial Engineering, 2010. **58**(2): p. 212-220.
43. Ronnberg, E. and T. Larsson, *Automating the self-scheduling process of nurses in Swedish healthcare: a pilot study*. Health Care Manag Sci, 2010. **13**(1): p. 35-53.
44. Topaloglu, S. and H. Selim, *Nurse scheduling using fuzzy modeling approach*. Fuzzy Sets and Systems, 2010. **161**(11): p. 1543-1563.
45. Brunner, J. and G. Edenharter, *Long term staff scheduling of physicians with different experience levels in hospitals using column generation*. Health Care Management Science, 2011. **14**(2): p. 189-202.
46. Chow, V.S., et al., *Reducing Surgical Ward Congestion Through Improved Surgical Scheduling and Uncapacitated Simulation*. Production and Operations Management, 2011. **20**(3): p. 418-430.
47. Mobasher, A., et al., *Daily scheduling of nurses in operating suites*. IIE Transactions on Healthcare Systems Engineering, 2011. **1**(4): p. 232-246.
48. Nilssen, E.J., et al., *Multi-skill shift design for norwegian hospitals*. Journal of Applied Operational Research, 2011. **3**(3): p. 137-147.
49. Topaloglu, S. and I. Ozkarahan, *A constraint programming-based solution approach for medical resident scheduling problems*. Computers & Operations Research, 2011. **38**(1): p. 246-255.
50. He, B., et al., *The Timing of Staffing Decisions in Hospital Operating Rooms: Incorporating Workload Heterogeneity into the Newsvendor Problem*. Manufacturing & Service Operations Management, 2012. **14**(1): p. 99-114.
51. He, F. and R. Qu, *A constraint programming based column generation approach to nurse rostering problems*. Computers & Operations Research, 2012. **39**(12): p. 3331-3343.
52. Valouxis, C., et al., *A systematic two phase approach for the nurse rostering problem*. European Journal of Operational Research, 2012. **219**(2): p. 425-433.
53. Yilmaz, E., *A Mathematical Programming Model for Scheduling of Nurses' Labor Shifts*. Journal of Medical Systems, 2012. **36**(2): p. 491-496.
54. Dohn, A. and A. Mason, *Branch-and-price for staff rostering: An efficient implementation using generic programming and nested column generation*. European Journal of Operational Research, 2013. **230**(1): p. 157-169.

55. Maenhout, B. and M. Vanhoucke, *An integrated nurse staffing and scheduling analysis for longer-term nursing staff allocation problems*. Omega, 2013. **41**(2): p. 485-499.
56. Sawik, B., *A single and triple-objective mathematical programming models for assignment of services in a healthcare institution*. International Journal of Logistics Systems and Management, 2013. **15**(2): p. 249-259.
57. Soto, R., et al., *Nurse and paramedic rostering with constraint programming: A case study*. Romanian Journal of Information Science and Technology, 2013. **16**(1): p. 52-64.
58. Todorovic, N. and S. Petrovic, *Bee Colony Optimization Algorithm for Nurse Rostering*. IEEE Transactions on Systems, Man, and Cybernetics: Systems, 2013. **43**(2): p. 467-473.
59. Baum, R., D. Bertsimas, and N. Kallus, *Scheduling, revenue management, and fairness in an academic-hospital radiology division*. Academic radiology, 2014. **21**(10): p. 1322-1330.
60. Harper, P.R., *A framework for operational modelling of hospital resources*. Health Care Manag Sci, 2002. **5**(3): p. 165-73.
61. Connelly, L.G. and A.E. Bair, *Discrete event simulation of emergency department activity: a platform for system-level operations research*. Acad Emerg Med, 2004. **11**(11): p. 1177-85.
62. Griffiths, J.D., et al., *Modelling the requirement for supplementary nurses in an intensive care unit*. Journal of the Operational Research Society, 2004. **56**(2): p. 126-133.
63. Marcon, E. and F. Dexter, *Impact of surgical sequencing on post anesthesia care unit staffing*. Health Care Management Science, 2006. **9**(1): p. 87-98.
64. Duguay, C. and F. Chetouane, *Modeling and Improving Emergency Department Systems using Discrete Event Simulation*. Simulation, 2007. **83**(4): p. 311-320.
65. Hung, G.R., et al., *Computer Modeling of Patient Flow in a Pediatric Emergency Department Using Discrete Event Simulation*. Pediatric Emergency Care, 2007. **23**(1): p. 5-10.
66. Molema, J.J., et al., *Healthcare system design and parttime working doctors*. Health Care Manag Sci, 2007. **10**(4): p. 365-71.
67. van Oostrum, J.M., et al., *A Simulation Model for Determining the Optimal Size of Emergency Teams on Call in the Operating Room at Night*. Anesthesia & Analgesia, 2008. **107**(5): p. 1655-1662 10.1213/ane.0b013e318184e919.
68. Sundaramoorthi, D., et al., *A data-integrated simulation model to evaluate nurse-patient assignments*. Health Care Manag Sci, 2009. **12**(252-268).
69. Wang, T., et al., *Modelling and simulation of emergency services with ARIS and Arena. Case study: the emergency department of Saint Joseph and Saint Luc Hospital*. Production Planning & Control, 2009. **20**(6): p. 484-495.
70. Brenner, S., et al., *Modeling and analysis of the emergency department at University of Kentucky Chandler Hospital using simulations*. Journal of Emergency Nursing, 2010. **36**(4): p. 303-310.
71. Day, T.E., et al., *The use of queueing and simulative analyses to improve an overwhelmed pharmacy call center*. J Pharm Pract, 2010. **23**(5): p. 492-5.

72. Harper, P.R., N.H. Powell, and J.E. Williams, *Modelling the size and skill-mix of hospital nursing teams*. The Journal of the Operational Research Society, 2010. **61**(5): p. 768-779.
73. Reynolds, M., et al., *Using discrete event simulation to design a more efficient hospital pharmacy for outpatients*. Health Care Manag Sci, 2011. **14**(3): p. 223-36.
74. Izady, N. and D. Worthington, *Setting staffing requirements for time dependent queueing networks: The case of accident and emergency departments*. European Journal of Operational Research, 2012. **219**(3): p. 531-540.
75. Azari-Rad, S., et al., *A simulation model for perioperative process improvement*. Operations Research for Health Care, 2014. **3**(1): p. 22-30.
76. Doan, Q., et al., *Forecasting the effect of physician assistants in a pediatric ED*. Journal of the American Academy of Physician Assistants, 2014. **27**(8): p. 35-41
10.1097/01.JAA.0000451860.95151.e1.
77. Kadri, F., S. Chaabane, and C. Tahon, *A simulation-based decision support system to prevent and predict strain situations in emergency department systems*. Simulation Modelling Practice and Theory, 2014. **42**: p. 32-52.
78. Ingalls, R.G. *Introduction to simulation*. in *Proceedings of the 40th Conference on Winter Simulation*. 2008. Winter Simulation Conference.
79. Brailsford, S., *Overcoming the barriers to implementation of operations research simulation models in healthcare*. Clinical and investigative medicine. Medecine clinique et experimentale, 2005. **28**(6): p. 312-315.
80. Eldabi, T., Z. Irani, and R.J. Paul, *A proposed approach for modelling health-care systems for understanding*. Journal of Management in Medicine, 2002. **16**(2/3): p. 170-187.
81. Mielczarek, B. and J. Uziółko-Mydlikowska, *Application of computer simulation modeling in the health care sector: a survey*. Simulation, 2010: p. 0037549710387802.
82. Harper, P. and M. Pitt, *On the challenges of healthcare modelling and a proposed project life cycle for successful implementation&star*. Journal of the Operational Research Society, 2004. **55**(6): p. 657-661.

Appendix A. Time Study Margin of Error Calculation

$$E = z_{\alpha/2} \frac{\sigma}{\sqrt{n}}$$

where


$z_{\alpha/2}$ = critical value of 95% confidence level = 1.96

n = sample size of each patient type

E = margin of error

σ = standard deviation of consult time (min)

Appendix B. Conceptual Model: Process Flow Map of PHC Patient Flow



Appendix C. Conceptual Model: PHC Patient, Facility, & Worker Assumptions

Category	Assumptions
<p style="text-align: center;">Patient Assumptions</p>	<ul style="list-style-type: none"> • Entities (patients) begin arriving at 7:30am • Patients inter-arrival time is exponentially distributed where the inter-arrival time is set at 1second <ul style="list-style-type: none"> ○ Assumes arrivals occur according to a homogenous Poisson process • Of outpatients, 70% of outpatients are adults, 30% are pediatric • 60% of Adult Outpatients require lab, pharmacy, or both services during a visit. Of these patients: <ul style="list-style-type: none"> ○ 60% require only pharmacy services ○ 10% require only lab services ○ 30% require both pharmacy and lab services • 45% of Pediatric Outpatients require lab, pharmacy, or both services during a visit. Of these patients: <ul style="list-style-type: none"> ○ 60% require only pharmacy services ○ 10% require only lab services ○ 30% require both pharmacy and lab services • 15% of Well Child Care Patients need to receive outpatient services. These 15% will receive the same services as a pediatric outpatient in addition to immunizations that occur during treatment by a nurse/doctor. • 10% of 1st Antenatal Care Patients require lab, pharmacy, or both services during a visit. Of these patients: <ul style="list-style-type: none"> ○ 60% require only pharmacy services ○ 10% require only lab services ○ 30% require both pharmacy and lab services • Seasonal factors do not affect average daily visits
<p style="text-align: center;">Facility Assumptions</p>	<ul style="list-style-type: none"> • Hours of operation at each healthcare facility are between 8:00am-2:00pm
<p style="text-align: center;">Worker Assumptions</p>	<ul style="list-style-type: none"> • Each resource (healthcare worker) provides services continuously during hours of operation

Appendix D. DES Model Validation Hypothesis Test

For each type of patient i and for each facility j
let X_{ij} = model output average delay time
 μ_{ij} = actual system average delay time

Hypothesis Test:

$$H_0 : E(X_{ij}) = \mu_{ij}$$

$$H_1 : E(X_{ij}) \neq \mu_{ij}$$

reject H_0 if $|t_0| > t_{\alpha/2, n-1}$

$$\text{Compute test statistic} = t_0 = \frac{\bar{X}_{ij} - \mu_{ij}}{s / \sqrt{n}}$$

where model sample mean = $\bar{X}_{ij} \approx E(X_{ij})$, $\bar{X}_{ij} = \frac{1}{n} \sum_{i=1}^n X_{ij}$

model sample standard deviation = $s = \frac{\sum_{i=1}^n (X_{ij} - \bar{X}_{ij})^2}{n - 1}$

Per a 2 – sided t – test and assuming $\alpha = 0.05$

critical value of $t = t_{\alpha/2, n-1} = t_{0.025, 4} = 2.776$

For each type of patient and for each facility

Appendix E. DES Model Validation Hypothesis Test Results Summary

Rural Hospital			Rural Health Center			Urban Health Center					
<i>Adult Outpatient</i>			<i>Adult Outpatient</i>			<i>Adult Outpatient</i>					
System Treatment Delay (min)	Model Treatment Delay (min)		System Treatment Delay (min)	Model Treatment Delay (min)		System Treatment Delay (min)	Model Treatment Delay (min)				
65		157.56	53		134.09	91		207			
63		142.49	30		137.59	42		195			
46		152.69	58		138.59	63		203			
75		143.40	54		130.00	65		197			
52		137.16	59		133.87	82		193			
Actual Average =	60.20	Model Average =	146.66	Actual Average =	50.80	Model Average =	134.83	Actual Average =	68.6	Model Average =	199.21
t ₀ =	23.38	Model St. Dev. =	8.27	t ₀ =	55.08	Model St. Dev. =	3.41	t ₀ =	50.88	Model St. Dev. =	5.74
Conclusion: Reject H₀			Conclusion: Reject H₀			Conclusion: Reject H₀					
<i>Pediatric Outpatient</i>			<i>Pediatric Outpatient</i>			<i>Pediatric Outpatient</i>					
System Treatment Delay (min)	Model Treatment Delay (min)		System Treatment Delay (min)	Model Treatment Delay (min)		System Treatment Delay (min)	Model Treatment Delay (min)				
18		98.05	67		98.05	45		182			
28		144.50	55		109.30	40		194			
36		140.76	46		97.58	38		196			
37		133.29	51		93.70	63		186			
20		151.40	64		108.56	46		186			
Actual Average =	27.80	Model Average =	133.60	Actual Average =	56.60	Model Average =	101.44	Actual Average =	46.4	Model Average =	188.81
t ₀ =	11.31	Model St. Dev. =	20.92	t ₀ =	14.22	Model St. Dev. =	7.05	t ₀ =	52.40	Model St. Dev. =	6.08
Conclusion: Reject H₀			Conclusion: Reject H₀			Conclusion: Reject H₀					
<i>1st Antenatal Care Patient</i>			<i>1st Antenatal Care Patient</i>			<i>1st Antenatal Care Patient</i>					
System Treatment Delay (min)	Model Treatment Delay (min)		System Treatment Delay (min)	Model Treatment Delay (min)		System Treatment Delay (min)	Model Treatment Delay (min)				
71		41.86	96		45.12	50		43			
77		36.02	99		48.46	78		40			
74		42.34	81		63.19	84		46			
42		38.52	86		46.98	9		41			
48		42.15	73		32.72	43		56			
Actual Average =	62.40	Model Average =	40.18	Actual Average =	87.00	Model Average =	47.29	Actual Average =	52.80	Model Average =	45.24
t ₀ =	-17.73	Model St. Dev. =	2.80	t ₀ =	-8.18	Model St. Dev. =	10.85	t ₀ =	-2.66	Model St. Dev. =	6.37
Conclusion: Reject H₀			Conclusion: Reject H₀			Conclusion: Fail to reject H₀					
<i>Immunization Patients</i>			<i>Immunization Patients</i>			<i>Immunization Patients</i>					
System Treatment Delay (min)	Model Treatment Delay (min)		System Treatment Delay (min)	Model Treatment Delay (min)		System Treatment Delay (min)	Model Treatment Delay (min)				
43		128.61	47		48.10	32		99.31			
38		147.04	37		44.62	29		101.24			
16		154.18	62		44.03	39		97.3347			
14		141.40	40		44.17	47		99.81			
9		144.63	62		44.19	41		103.27			
Actual Average =	24.00	Model Average =	143.17	Actual Average =	49.60	Model Average =	45.02	Actual Average =	37.60	Model Average =	100.19
t ₀ =	28.35	Model St. Dev. =	9.40	t ₀ =	-5.89	Model St. Dev. =	1.74	t ₀ =	63.13	Model St. Dev. =	2.22
Conclusion: Reject H₀			Conclusion: Reject H₀			Conclusion: Reject H₀					

Appendix F. Rural Health Center Orthogonal Array for What-If Scenario Experiments

Rural Health Center Orthogonal Array for What-If Scenario Experiments					
Exp #	Registration Agent	ANC RN	Pharm Worker	Lab Worker	Immunization Tech
1	2	2	1	1	1
2	1	1	4	4	2
3	3	6	5	6	4
4	4	5	4	1	3
5	2	4	2	6	2
6	1	6	1	2	1
7	5	1	1	1	3
8	2	5	1	5	5
9	2	5	2	4	6
10	5	3	5	1	6
11	2	2	6	4	2
12	3	3	6	4	3
13	3	1	5	5	1
14	3	2	4	5	6
15	6	5	3	1	1
16	4	5	6	5	2
17	1	6	6	3	6
18	2	1	3	6	3
19	4	2	1	4	5
20	6	5	4	2	6
21	5	2	4	6	4
22	4	3	6	6	6
23	5	5	5	3	2
24	2	2	5	1	1
25	1	5	2	6	3
26	3	5	3	3	5
27	3	1	3	4	6
28	4	1	2	5	4
29	4	6	3	4	1
30	5	5	6	2	4
31	4	1	5	6	5
32	6	6	3	6	2
33	4	3	5	2	3
34	5	6	2	5	5
35	6	1	2	2	2
36	5	6	4	4	3
37	1	4	1	5	4
38	4	4	3	3	4
39	1	2	3	5	3
40	1	2	5	2	2
41	6	4	6	5	3
42	6	6	5	5	6
43	3	4	6	2	5
44	6	1	4	3	4
45	6	3	1	4	4
46	4	6	1	1	2
47	1	1	6	1	5
48	2	3	4	5	1
49	6	3	2	1	5
50	5	3	3	5	2
51	4	4	4	2	1
52	3	5	1	6	1
53	3	2	2	1	4
54	1	3	4	6	5
55	2	6	4	3	5
56	2	6	6	1	4
57	5	1	6	3	1
58	3	4	4	1	2
59	3	3	1	3	2
60	1	4	3	1	6
61	1	3	2	3	1
62	3	6	2	2	3
63	6	4	5	4	5
64	5	2	3	2	5
65	2	4	5	3	3
66	2	1	1	2	6
67	5	4	1	6	6
68	2	3	3	2	4
69	4	2	2	3	6
70	6	2	1	3	3
71	6	2	6	6	1
72	1	5	5	4	4
73	5	4	2	4	1


Appendix G. Urban Health Center Orthogonal Array for What-If Scenario Experiments

Urban Health Center Orthogonal Array for What-If Scenario Experiments							
Exp #	Registration Agent	Outpatient RN	Outpatient MD	ANC RN	Pharm Worker	Lab Worker	Immunization Tech
1	1	9	1	5	6	4	2
2	2	10	3	7	9	7	5
3	5	8	3	7	4	3	1
4	2	8	6	4	6	3	4
5	5	11	4	6	8	6	5
6	4	11	6	3	5	7	2
7	3	9	2	4	5	7	4
8	5	7	5	5	5	2	4
9	1	11	1	5	9	7	1
10	2	9	4	6	5	5	2
11	5	11	2	3	9	4	4
12	3	11	3	8	7	3	6
13	6	10	2	5	8	3	2
14	4	10	1	4	4	6	6
15	1	7	4	3	6	3	2
16	4	7	2	6	7	3	3
17	3	12	4	4	8	7	3
18	6	12	6	6	7	7	1
19	2	7	3	4	9	6	3
20	5	9	6	5	4	3	5
21	1	12	3	5	6	6	2
22	2	11	2	7	4	5	2
23	6	7	1	8	4	7	2
24	4	8	4	8	9	2	5
25	3	8	1	5	6	5	3
26	2	8	5	3	7	7	5
27	1	8	2	4	8	2	6
28	6	12	2	5	7	2	5
29	6	11	5	4	6	4	1
30	6	9	4	7	9	3	3
31	4	8	4	5	4	7	4
32	3	10	5	6	9	2	2
33	5	12	1	7	5	2	3
34	4	12	5	3	9	3	6
35	6	7	5	4	4	5	5
36	5	10	4	4	7	5	1
37	1	9	5	6	4	4	3
38	2	11	5	5	7	6	3
39	3	8	1	7	7	4	2
40	2	10	4	5	5	4	6
41	2	12	1	6	8	3	4
42	6	9	1	8	9	6	4
43	3	10	5	8	5	3	1
44	3	11	3	6	4	2	4
45	1	11	1	4	5	3	5
46	4	10	1	6	6	4	5
47	1	9	5	7	8	7	6
48	5	10	2	8	6	7	3
49	5	9	1	3	7	5	6
50	4	11	6	8	8	5	3
51	2	12	2	8	4	4	6
52	4	9	3	4	7	2	2
53	2	7	1	3	8	2	1
54	6	8	6	6	5	6	6
55	3	12	4	3	4	6	1
56	4	9	3	5	8	4	1
57	6	10	3	3	8	5	4
58	6	11	4	7	6	2	6
59	5	12	6	4	9	4	2
60	6	8	3	3	5	4	3
61	3	9	2	3	6	6	5
62	1	7	4	8	7	4	4
63	1	10	6	7	7	6	4
64	3	7	6	7	8	4	5
65	5	7	3	6	6	7	6
66	5	8	5	8	8	6	2
67	1	10	6	3	4	2	3
68	4	12	5	7	6	5	4
69	3	7	6	5	9	5	6
70	4	7	2	7	5	6	1
71	2	9	6	8	6	2	1
72	1	12	3	8	5	5	5
73	1	8	2	6	9	5	1

Appendix H. Rural Hospital Orthogonal Array for What-If Scenario Experiments

Rural Hospital Orthogonal Array for What-If Scenario Experiments								
Exp #	Registration Agent (Primary)	Registration Agent (Secondary)	Outpatient RN	Outpatient MD	ANC RN	Pharm Worker	Lab Worker	Immunization Tech
1	2	1	14	1	5	3	2	2
2	2	4	16	3	3	2	4	5
3	6	6	12	4	6	3	1	2
4	2	4	12	1	5	6	1	6
5	3	1	15	1	8	2	6	4
6	1	6	16	3	5	6	5	1
7	4	2	16	1	7	4	2	1
8	3	6	14	6	7	2	5	5
9	6	1	14	2	3	3	5	6
10	3	5	17	4	5	4	5	6
11	5	5	15	1	5	1	3	3
12	1	5	14	1	4	5	4	4
13	3	6	15	3	6	6	4	6
14	5	1	13	5	7	6	2	6
15	2	3	14	5	7	1	6	3
16	6	1	15	5	5	4	4	1
17	3	4	13	2	6	3	4	3
18	2	3	15	4	6	2	3	2
19	1	1	13	3	6	5	2	1
20	1	5	17	3	6	4	5	2
21	3	1	17	6	3	6	3	3
22	6	5	12	2	6	1	2	6
23	6	6	16	5	7	5	4	3
24	5	4	14	6	4	1	4	6
25	4	6	14	5	4	6	3	2
26	2	3	17	2	4	6	2	4
27	4	5	13	4	5	2	5	3
28	5	4	17	1	6	6	5	5
29	2	1	13	4	8	6	1	4
30	5	6	16	6	8	3	3	4
31	4	3	12	1	3	2	5	1
32	5	5	17	6	8	2	2	2
33	3	1	14	3	5	1	2	2
34	1	2	13	4	3	1	4	2
35	6	4	13	1	3	4	3	4
36	4	2	12	6	6	5	6	5
37	2	6	17	5	8	4	6	6
38	6	3	17	1	7	2	4	6
39	4	3	13	6	5	6	4	5
40	3	6	17	1	4	1	1	1
41	5	2	15	5	8	5	5	6
42	2	1	12	5	5	2	4	2
43	2	5	13	3	7	4	6	5
44	3	5	15	5	3	3	1	5
45	4	5	16	2	8	6	4	2
46	5	3	16	5	6	3	5	4
47	4	1	14	1	7	3	5	2
48	4	1	17	3	4	5	1	3
49	5	1	12	2	4	2	3	5
50	5	2	14	2	5	4	1	5
51	2	2	14	1	6	4	1	3
52	6	5	16	4	7	6	3	5
53	1	2	12	5	8	6	5	3
54	2	2	15	6	4	5	5	2
55	2	2	17	3	7	3	4	4
56	5	1	16	4	6	1	6	1
57	1	4	14	5	6	2	2	4
58	6	4	16	6	5	5	2	3
59	1	3	15	2	8	4	3	1
60	3	5	14	2	8	5	4	1
61	6	2	15	1	3	6	2	2
62	1	6	13	6	3	2	1	6
63	6	2	17	6	5	1	6	4
64	2	6	14	4	3	5	3	1
65	2	1	16	2	3	1	5	3
66	6	3	15	3	4	1	5	5
67	1	6	12	1	8	1	4	5
68	5	4	15	3	7	2	1	1
69	3	2	13	5	6	1	3	5
70	3	2	12	2	7	6	6	1
71	3	3	12	3	5	5	3	4
72	5	6	12	3	3	4	2	3
73	4	4	15	2	3	5	6	6
74	5	2	17	4	3	3	4	1
75	6	2	14	3	8	2	3	3
76	1	4	17	2	7	1	3	2
77	3	3	13	1	8	3	2	3
78	4	5	12	5	3	1	1	4
79	1	3	17	5	3	5	2	5
80	1	1	16	1	4	3	6	5
81	3	4	16	5	4	4	1	2
82	3	4	12	4	7	5	5	4
83	4	2	13	3	4	3	3	6
84	6	1	17	4	8	5	1	5
85	2	4	13	6	8	1	5	1
86	5	6	13	1	5	5	6	2
87	6	3	14	6	6	6	1	1
88	4	6	15	4	7	1	2	4
89	4	1	15	6	6	4	4	4
90	4	6	17	2	6	2	6	3
91	1	2	16	2	5	2	1	4
92	4	4	14	4	8	4	2	5
93	1	3	14	4	5	3	6	6
94	2	6	15	2	5	3	2	5
95	2	5	12	6	4	3	2	1
96	5	5	14	3	3	6	6	4
97	1	1	12	6	7	4	3	6
98	4	3	16	3	8	1	1	6
99	5	3	13	2	7	5	1	2
100	6	6	13	2	4	4	5	4
101	3	2	16	4	4	2	2	6
102	2	5	16	1	6	5	3	6
103	5	3	12	4	4	4	4	3
104	1	4	15	4	4	6	6	3
105	4	4	17	5	5	3	3	1
106	6	5	13	5	4	2	6	1
107	6	4	12	3	8	3	6	2
108	1	5	15	6	7	3	1	3
109	3	3	16	6	3	4	6	2

Appendix I. Spreadsheet-based Decision Support Tool Example



Moz DSS3 [Read-Only] - Microsoft Excel (Product Activation Failed)

File Home Insert Page Layout Formulas Data Review View Developer

Calibri 14 A A

General

E9 60

A B C D E F G H I

1 *Clique* nas caixas brancas abaixo e *digite* o numero de visitas diarias esperados tanto para adulto ambulatorial como para pacientes pediatricas.
 2 Pressione *Enter* para ver o numero de profissionais de saude requerido
 3 Pressione *Reset* para restabelecer esta tela
 4

Adult Oupatient Visits per Day	Pediatric Patient Visits per Day	Well Child Patient Visits per Day	ANC Patient Visits per Day
40	22	34	12

**Target Total
Average Wait
Time**

60

BACK RESET ENTER

Facility InterfaceRHC

Moz DSS3 [Read-Only] - Microsoft Excel (Product Activation Failed)

File Home Insert Page Layout Formulas Data Review View Developer

Calibri 14 A A

General

Pharmacy Worker

1 O seguinte sao os numeros de profissionais de saude necessarios de modo que
 2 os pacientes tenham um tempo de espera total media inferior a 1 hora
 3

Outpatient RNs	4	Immunization Techs	6
ANC RNs	3	Laboratory Worker	6
		Pharmacy Worker	6

BACK END

InterfaceRHC EndRHC

Appendix J. Regression Models and Statistical Summaries

Rural Health Center						
Response Variable	Model	Lack-of-fit	NOVA F Tes	R ² adj.	Durbin-Watson Statistic	Anderson Darlin [p-value]
Adult Outpatient Wait Time	AWT = 120.667 + 1.10595 AV - 19.1836 BRNV - 0.696486 LabV - 0.557691 PharmV	P>0.05	p<0.05	83.49%	1.73	25.12 [p<0.005]
Adult Outpatient Wait Time [BE]	AWT = 118.814 + 1.106 AV - 19.1923 BRNV - 0.711789 LabV	p<0.05	p<0.05	83.47%	1.73	24.60 [p<0.005]
Transformed Adult Outpatient Wait Time	AWT*0.227453 = 2.8627 + 0.00718114 AV - 0.10564 BRNV - 0.00453417 LabV - 0.00397788 PharmV	P>0.05	p<0.05	86.78%	1.09	22.72 [p<0.005]
Pediatric Outpatient Wait Time	PWT = 94.0721 + 1.94099 PV - 14.6733 BRNV - 0.344633 LabV - 0.267091 PharmV	P>0.05	p<0.05	75.68%	1.91	47.39 [p<0.005]
Pediatric Outpatient Wait Time [BE]	PWT = 91.9837 + 1.94109 PV - 14.6833 BRNV	p<0.05	p<0.05	75.67%	1.9	46.34 [p<0.005]
Transformed Ped. Outpatient Wait Time	PWT*-0.0680245 = -0.744201 + 0.00119551 PV - 0.00731516 BRNV - 0.000272647 LabV - 0.000243344 PharmV	P>0.05	p<0.05	83.95%	1.10	20.60 [p<0.005]
Transformed Ped. Outpatient Wait Time [BE]	PWT*-0.0673181 = -0.747284 + 0.00118675 PV - 0.00726617 BRNV - 0.000272242 LabV	p<0.05	p<0.05	83.92%	1.10	20.26 [p<0.005]
Well Child Care Patient	WCWT = 40.2754 + 0.453931 WCV - 0.914471 BRNV - 0.0599932 LabV - 0.0562125 PharmV - 1.97145 PreTV	P>0.05	p<0.05	78.65%	2.04	33.43 [p<0.005]
Well Child Care Patient [BE]	WCWT = 39.8895 + 0.453956 WCV - 1.97461 PreTV - 0.916315 BRNV	p<0.05	p<0.05	78.63%	2.04	33.77 [p<0.005]
Transformed Well Child Care Patient	(WCWT*-1)/(L*g*(L-1)) = 146085 + 0.434451 WCV - 1.47802 PreTV - 0.850668 BRNV - 0.0012703 PharmV - 0.033537 LabV L = Lambda = -2.53882, g = 37.4774 is the geometric mean of WCWT	P>0.05	p<0.05	88.63%	1.61	3.85 [p<0.005]
Transformed Well Child Care Patient [BE]	(WCWT*-1)/(L*g*(L-1)) = 146085 + 0.434457 WCV - 1.47897 PreTV - 0.851227 BRNV L = Lambda = -2.53882, g = 37.4774 is the geometric mean of WCWT	p<0.05	p<0.05	88.63%	1.61	4.09 [p<0.005]
Pregnant Patient Wait Time	PregWT = 38.6559 + 1.48583 PRV - 2.43632 ERNV - 0.0481731 PharmV - 0.036288 LabV	p<0.05	p<0.05	62.97%	2.05	40.24 [p<0.005]
Pregnant Patient Wait Time [BE]	PregWT = 38.3682 + 1.48592 PRV - 2.43789 ERNV	p<0.05	p<0.05	63.00%	2.05	40.19 [p<0.005]
Transformed Preg. Patient Wait Time	(PregWT*-1)/(L*g*(L-1)) = 535023 + 1.26618 PRV - 1.64813 ERNV - 0.0606347 PharmV - 0.0654934 LabV L = Lambda = -3, g = 35.5939 is the geometric mean of PregWT	p<0.05	p<0.05	70.43%	1.84	4.04 [p<0.005]
Transformed Preg. Patient Wait Time [BE]	(PregWT*-1)/(L*g*(L-1)) = 535022 + 1.26631 PRV - 1.65048 ERNV L = Lambda = -3, g = 35.5939 is the geometric mean of PregWT	p<0.05	p<0.05	70.37%	1.84	3.86 [p<0.005]

Urban Health Center						
Response Variable	Model	Lack-of-fit		R ² adj.	Durbin-Watson Statistic	Anderson Darlin [p-value]
Adult Outpatient Wait Time	AWT = 128.727 + 0.121836 AV - 0.0843473 RegV - 3.91733 BRNV - 3.90924 MDV - 0.376547 PharmV - 0.135669 LabV	p>0.05	p<0.05	84.88%	0.29	34.68 [p<0.005]
Adult Outpatient Wait Time [BE]	AWT = 126.619 + 0.121836 AV - 3.91808 BRNV - 3.91301 MDV	p<0.05	p<0.05	84.89%	0.30	33.95 [p<0.005]
Transformed Adult Outpatient Wait Time	(AWT*-1)/(L*g*(L-1)) = 61.62 + 0.118828 AV - 0.0756183 RegV - 4.49829 BRNV - 4.5089 MDV - 0.481731 PharmV + 0.495465 LabV L = Lambda = 2.58286, g = 146.130 is the geometric mean of AWT	p>0.05	p<0.05	91.52%	0.93	8.74 [p<0.005]
Transformed Adult Outpatient Wait Time [BE]	(AWT*-1)/(L*g*(L-1)) = 61.3739 + 0.118828 AV - 4.49873 BRNV - 4.51107 MDV - 0.482164 PharmV + 0.49504 LabV L = Lambda = 2.58286, g = 146.130 is the geometric mean of AWT	p>0.05	p<0.05	91.52%	0.93	8.71 [p<0.005]
Pediatric Outpatient Wait Time	PWT = 125.118 + 0.293655 PV - 0.0920016 RegV - 4.19844 BRNV - 4.20444 MDV - 0.265244 PharmV + 0.128185 LabV	p>0.05	p<0.05	87.50%	0.35	24.73 [p<0.005]
Pediatric Outpatient Wait Time [BE]	PWT = 123.671 + 0.293655 PV - 4.19912 BRNV - 4.20784 MDV	p<0.05	p<0.05	87.52%	0.35	24.17 [p<0.005]
Transformed Ped. Outpatient Wait Time	(PWT*-1)/(L*g*(L-1)) = 65.3777 + 0.28858 PV - 0.0823304 RegV - 4.69906 BRNV - 4.70601 MDV - 0.322257 PharmV - 0.340038 LabV L = Lambda = 2.25076, g = 142.067 is the geometric mean of PWT	p>0.05	p<0.05	91.80%	0.90	8.29 [p<0.005]
Transformed Ped. Outpatient Wait Time [BE]	(PWT*-1)/(L*g*(L-1)) = 64.5921 + 0.288538 PV - 4.69722 BRNV - 4.70596 MDV L = Lambda = 2.24704, g = 142.067 is the geometric mean of PWT	p<0.05	p<0.05	91.78%	0.91	7.21 [p<0.005]
Well Child Care Patient	WCWT = 94.0701 + 0.446264 WCV - 3.75518 RegV - 0.764232 BRNV + 0.295787 MDV - 12.231 PreTV + 0.161791 PharmV - 0.745296 LabV	p>0.05	p<0.05	77.32%	1.96	54.56 [p<0.005]
Well Child Care Patient [BE]	WCWT = 96.0686 + 0.446264 WCV - 3.74594 RegV - 0.762384 BRNV - 12.2255 PreTV - 0.743447 LabV	p>0.05	p<0.05	77.33%	1.96	55.44 [p<0.005]
Transformed Well Child Care Patient	WCWT*-0.340488 = -0.236239 + 0.000483037 WCV - 0.00302548 RegV - 0.0103864 PreTV - 0.000474143 BRNV + 0.000335128 MDV - 0.000129612 PharmV - 0.000655445 LabV	p>0.05	p<0.05	89.92%	1.13	8.33 [p<0.005]
Transformed Well Child Care Patient [BE]	WCWT*-0.340429 = -0.23605 + 0.000483073 WCV - 0.00301704 RegV - 0.010382 PreTV - 0.000472447 BRNV - 0.000653759 LabV	p>0.05	p<0.05	89.91%	1.13	8.60 [p<0.005]
Pregnant Patient Wait Time	PregWT = 47.8384 + 0.829958 PRV - 3.26423 ERNV - 0.0456434 PharmV - 0.361975 LabV	p>0.05	p<0.05	88.49%	1.91	30.26 [p<0.005]
Pregnant Patient Wait Time [BE]	PregWT = 47.5426 + 0.829958 PRV - 3.26429 ERNV - 0.362029 LabV	p<0.05	p<0.05	88.49%	1.91	30.07 [p<0.005]
Transformed Preg. Patient Wait Time	PregWT*-0.713901 = -0.0685864 + 0.000905954 PRV - 0.00296484 ERNV - 5.83559e-005 PharmV - 0.000390646 LabV	p>0.05	p<0.05	93.51%	1.35	6.59 [p<0.005]
Transformed Preg. Patient Wait Time [BE]	PregWT*-0.713506 = -0.0690655 + 0.000906793 PRV - 0.00296778 ERNV - 0.000391078 LabV	p<0.05	p<0.05	93.51%	1.35	6.52 [p<0.005]

Rural Hospital						
Response Variable	Model	Lack-of-fit		R ² adj.	Durbin-Watson Statistic	Anderson Darlin [p-value]
Adult Outpatient Wait Time	AWT = 143.52 + 0.325805 AV - 4.48906 RegV1 - 3.55044 BRNV - 3.43398 MDV - 2.02939 PharmV - 0.53778 LabV	p>0.05	p<0.05	90.86%	1.39	11.90 [p<0.005]
Transformed Adult Wait Time	AWT*1.26787 = 573.101 + 1.51659 AV - 22.1621 RegV1 - 17.415 BRNV - 16.9099 MDV - 9.43802 PharmV - 1.95441 LabV	p>0.05	p<0.05	91.35%	1.73	6.00 [p<0.005]
Pediatric Outpatient Wait Time	PWT = 84.2908 + 0.617292 PV + 10.0146 RegV1 - 3.38159 BRNV - 3.85698 MDV - 2.34554 PharmV - 1.29786 LabV	p>0.05	p<0.05	80.91%	1.99	19.03 [p<0.005]
Transformed Ped. Outpatient Wait Time	PWT*0.433749 = 6.5857 + 0.0193667 PV + 0.286226 RegV1 - 0.0945083 BRNV - 0.103485 MDV - 0.0703437 PharmV - 0.0487926 LabV	p>0.05	p<0.05	83.15%	1.56	6.16 [p<0.005]
Well Child Care Patient	WCWT = 115.839 + 0.490139 WCV - 1.23523 RegV2 - 18.1671 PreTV - 0.0863572 BRNV - 0.489667 MDV - 0.0303029 PharmV - 0.525024 LabV	p>0.05	p<0.05	88.17%	1.80	19.48 [p<0.005]
Well Child Care Patient [BE]	WCWT = 114.486 + 0.490139 WCV - 1.23567 RegV2 - 18.1674 PreTV - 0.490103 MDV - 0.525285 LabV	p>0.05	p<0.05	88.18%	1.80	19.33 [p<0.005]
Transformed Well Child Care Patient	WCWT*0.397467 = 6.2631 + 0.0125831 WCV - 0.0256401 RegV2 - 0.420784 PreTV - 0.00230074 BRNV - 0.0110497 MDV + 0.000265971 PharmV - 0.0137362 LabV	p>0.05	p<0.05	89.64%	1.35	28.42 [p<0.005]
Transformed Well Child Care Patient [BE]	WCWT*0.397497 = 6.23164 + 0.0125857 WCV - 0.0256534 RegV2 - 0.420879 PreTV - 0.0110597 MDV - 0.0137437 LabV	p>0.05	p<0.05	89.65%	1.35	28.40 [p<0.005]
Pregnant Patient Wait Time	PregWT = 45.5047 + 0.962978 PRV - 2.61967 ERNV - 0.207573 PharmV - 0.3907 LabV	p<0.05	p<0.05	88.36%	1.68	27.21 [p<0.005]
Transformed Preg. Patient Wait Time	PregWT*-0.218378 = -0.44138 + 0.00217199 PRV - 0.00516606 ERNV - 0.000414973 PharmV - 0.000834499 LabV	p<0.05	p<0.05	90.69%	1.44	7.10 [p<0.005]

BE=Backwards Elimination: the resulting model and statistic after independent variables with a p-value >0.05 were removed and the regression was performed again
Selected for decision support tool