

Lupus IgA1 autoantibodies synergize with IgG to enhance pDC responses to
RNA-containing immune complexes

Hayley Ryan Waterman

A dissertation

submitted in partial fulfillment of the
requirements for the degree of

Doctor of Philosophy

University of Washington

2023

Reading Committee:

Jessica A. Hamerman, Chair

Adam Lacy-Hulbert

Ram Savan

Program Authorized to Offer Degree:

Molecular and Cellular Biology

© Copyright 2023

Hayley Ryan Waterman

University of Washington

Abstract

Lupus IgA1 autoantibodies synergize with IgG to enhance pDC responses to
RNA-containing immune complexes

Hayley Ryan Waterman

Chair of the Supervisory Committee

Jessica A. Hamerman

Department of Immunology

Autoantibodies to nuclear antigens are hallmarks of the autoimmune disease systemic lupus erythematosus (SLE) where they contribute to pathogenesis. However, there remains a gap in our knowledge regarding how different isotypes of autoantibodies contribute to disease, including the production of the critical type I interferon (IFN) cytokines by plasmacytoid dendritic cells (pDCs) in response to immune complexes (ICs). We focused on IgA, which is the second most prevalent isotype in serum, and along with IgG is deposited in glomeruli in lupus nephritis. Here, we show that individuals with SLE have IgA autoantibodies against most nuclear antigens, correlating with IgG against the same antigen. We investigated whether IgA autoantibodies against a major SLE autoantigen, Smith ribonucleoproteins (Sm/RNPs), play a role in IC activation of pDCs. We found that pDCs express the IgA-specific Fc receptor, Fc α R, and there was a striking ability of IgA1 autoantibodies to synergize with IgG in RNA-containing

ICs to generate robust pDC IFN α responses. pDC responses to these ICs required both Fc α R and Fc γ RIIa, showing a potent synergy between these Fc receptors. Sm/RNP IC binding to and internalization by pDCs were greater when ICs contained both IgA1 and IgG. In addition, binding of Sm/RNP ICs generated with IgA1-sufficient serum correlated to pDC Fc α R expression, but not Fc γ RIIa expression. Lastly, pDCs from individuals with SLE had higher binding of IgA1-containing ICs and higher expression of Fc α R than pDCs from healthy control individuals. Taken together, our data show a new mechanism by which IgA1 anti-nuclear antibodies contribute to SLE pathogenesis.

Table of contents

Title page	
Copy right information	
Abstract	
Table of contents	i
List of figures	ii
List of abbreviations	v
Acknowledgements	vi
Funding	vii
Background	1
Systemic lupus erythematosus is a multifaceted autoimmune disease	1
Antinuclear antibodies are a commonality in SLE	1
Plasmacytoid dendritic cells and type I IFNs are implicated in SLE pathogenesis	2
Plasmacytoid dendritic cells release type I IFNs in response to immune complexes	3
SLE studies have mainly focused on IgG isotype autoantibodies	3
Evidence that IgA autoantibodies could be relevant in SLE	4
IgA and the IgA specific receptor, Fc α R	5
Results	7
Introduction	7
SLE donors have IgA isotype ANAs to multiple nuclear autoantigens	8
pDCs express the IgA-specific Fc α R and bind IgA	9
IgA1 in Sm/RNP immune complexes enhances pDC IFN α responses via Fc α R and Fc γ RIIa co-engagement	10
IgA1 in immune complexes promotes internalization by pDCs	13

SLE pDCs have increased capacity to bind ICs and increased surface Fc α R expression compared to healthy control pDCs	15
Main Figures and Table	17
Supplementary Figures and Tables	29
Closing remarks	47
Discussion	47
Limitations	52
Outstanding questions and future directions	53
Methods	56
Human samples	56
IgG and IgA Nuclear Antigen Microarray	56
PBMC isolation and pDC enrichment from whole blood	57
Staining for Fc α R and Fc γ RIIa	57
Staining for IgA and IgG	58
Anti-Sm/RNP and total antibody ELISAs	58
IgA/IgG IC ELISA	59
IgA1 depletion from serum	60
Generation of Sm/RNP immune complexes	60
Immune complex stimulation of pDCs	61
Cytokine measurements from supernatants	61
Interferon Stimulated Genes (ISGs) measurement from whole blood	62
Immune complex internalization microscopy	63
Human kidney biopsy	63
Statistics	64
References	65

List of Figures

Main Figures	17
Figure 1. IgA and IgG RNA-associated anti-nuclear antibodies in SLE serum	17
Figure 2. Human blood pDCs express Fc α R and Fc γ RIIa and bind IgA and IgG	19
Figure 3. RNA-containing immune complexes generated from SLE donors with anti-Sm/RNP antibodies	21
Figure 4. IgA1 in Sm/RNP immune complexes enhances IFN α production by human pDCs	22
Figure 5: IgA1 autoantibodies contribute to immune complex association with pDCs	24
Figure 6: IgA1 autoantibodies enhance pDC internalization of immune complexes	26
Figure 7: SLE pDCs have increased binding to immune complexes and increased expression of Fc α R	27
Table 1: SLE Serum Donor Participants Summary	28

Supplementary Figures	29
Supplementary Figure 1: Supplementary figures relating to main Figure 1	29
Supplementary Figure 2: Supplementary figures relating to main Figure 2	31
Supplementary Figure 3: Supplementary figures relating to main Figure 3	33
Supplementary Figure 4: Supplementary figures relating to main Figure 4	35
Supplementary Figure 5: Supplementary figures relating to main Figure 5	37
Supplementary Figure 6: Representative images used to distinguish Sm/RNP-AF647 SLE serum immune complex internalization versus binding to pDCs	38
Supplementary Figure 7: Supplementary figures relating to main Figure 7	40
Supplementary Figure 8: Top 50 genes positively correlated with Fc α R MFI from whole blood RNA-Seq	41
Supplementary Table 1: SLE and HC Participant Population Summary	43
Supplementary Table 2: SLE Participant Detailed Summary	44
Supplementary Table 3: SLE and HC Participant Population Summary	45
Supplementary Table 4: Flow Cytometry Antibody Summary	46

List of abbreviations (alphabetically):

ANA – Anti-nuclear antibody

dsDNA – Double stranded DNA

ELISA – Enzyme linked immunosorbent assay

HC – Healthy Control

IC – Immune complex

IFN – Interferon

IgA – Immunoglobulin A

IgD – Immunoglobulin D

IgG – Immunoglobulin G

IgE – Immunoglobulin E

IgM – Immunoglobulin M

ISG – Interferon stimulated gene

ITAM - immune tyrosine activating motif

Fab – Fragment antigen binding

Fc – Fragment crystallizable

FcR – Fragment crystallizable receptor

pDC – Plasmacytoid dendritic cell

RNP – Ribonucleoprotein

SLE – Systemic lupus erythematosus

Sm – Smith (protein)

Sm/RNPs – Smith (protein)/ribonucleoprotein

TLR – Toll-like receptor

Acknowledgements

I would like to thank my mentor, Jessica Hamerman who supported me and my ideas and remained a constant source of encouragement and optimism throughout my project. She is also the only reason there are not many more missing or misplaced commas in this manuscript. I would also like to thank all the other Hamerman lab members of past and present for making the lab such a great place to work and for listening to me talk about whatever hobby, movie, tv show or book series was my current hyperfocus. I am grateful for everyone at BRI who I have gotten to know over the years and who have been involved in my project. I am also thankful to the MCB support staff for promptly responding all my questions and cheering me on. I would also like to thank all the human subject blood donors who made my work possible. I would like to thank my committee members for their help and guidance. And I would like to thank my first mentor, Jim Zimring, who gave me my first job out of undergraduate school, taught me the importance of controls and inspired me to progress my scientific career.

I would like to thank all my friends and family who do not always understand what it is that I do but are always so encouraging. And I would like to thank my friends in science who do mostly understand my research and have listened to me talk about my project over the years. I am also thankful to my mom whose love and support helped me be a curious kid and functioning adult. And my grandma who gave me the Eyewitness books that first made me interested in science.

Lastly, I would like to thank my wife, Rachel whose love and emotional and financial support were critical to my success as a graduate student. And my baby Charley, who undoubtedly made my last ten months of graduate school harder than not having a baby but who also slept through the night most of that time. I love you both so much.

Funding

This work was supported by National Institute of Health grants ITHS TL1 TR002318 to H.R.W., R21AI154841 to J.A.H., and R01AR076242 to J.A.H, and by the Lupus Research Alliance Lupus Targets and Mechanisms Award to J.A.H.

Background

Systemic lupus erythematosus is a multifaceted autoimmune disease

Systemic lupus erythematosus (SLE) is a complex chronic autoimmune disease characterized by the breakdown of tolerance to nuclear antigens, and dysregulated cytokine production. SLE is a lifelong disease that can affect a variety of organs including the skin, joints, kidney, lung, brain, gastrointestinal tract, and cardiovascular system. There is major variation in the combinations and severity of symptoms that present in each individual patient, making SLE especially difficult to diagnose correctly. As with other systemic autoimmune diseases, there is a strong sex bias in SLE, with a ratio of female to male patients of nine to one¹. Sex hormones alter immune phenotypes in humans and have been shown to contribute directly to SLE in mouse models¹⁻³. Additionally, *TLR7* dosage is a strong factor contributing to SLE pathogenesis and is located on the X chromosome where it can escape X chromosome inactivation in immune cells⁴. In the United States SLE is one of the leading causes of death in young females and is more prevalent with even higher morbidity and mortality rates for SLE patients with Black, Hispanic, or Asian ancestries^{5,6}. The heterogeneity of disease manifestations and our incomplete understanding of pathogenesis makes SLE a difficult disease to identify and treat effectively. Therapeutics for individuals with SLE have mainly relied on non-specific immune suppressive drugs such as steroids and hydroxychloroquine, and only recently have more targeted therapies been approved for SLE⁷⁻⁹. Thus, there is a need for better understanding disease mechanisms in SLE for the development of new therapies and better diagnosis.

Antinuclear antibodies are a commonality in SLE

Despite the variety in disease manifestations, a hallmark of SLE is the breakdown of B cell tolerance to nuclear antigens that results in the presence of circulating autoreactive anti-nuclear

antibodies (ANAs). ANAs recognize nuclear antigens such as DNA or RNA themselves, or DNA or RNA-associated proteins, such as histones or subunits of ribonucleoproteins (RNPs)¹⁰⁻¹². The titers of some ANA specificities have associations with SLE disease activity and particular manifestations. For example, anti-nucleosome IgG titers are correlated to disease activity and lupus nephritis¹³. ANAs form nucleic acid-containing immune complexes (ICs) which deposit in tissues where they are recognized by and trigger inflammatory responses in Fc receptor (FcR) expressing cells such as dendritic cells, monocytes, and neutrophils¹⁴⁻¹⁸. In this way ANA-ICs facilitate nucleic acid entry to endosomes where RNA and DNA sensing innate immune receptors reside, thereby allowing immune cells to aberrantly respond to autologous RNA and DNA molecules.

Plasmacytoid dendritic cells and type I IFNs are implicated in SLE pathogenesis

Plasmacytoid dendritic cells (pDCs) are specialized innate immune cells that release large amounts of type I interferon (IFN) cytokines in response to RNA and DNA sensing by toll-like receptors (TLRs) 7 and 9, respectively¹⁸⁻²². pDCs can make up to 1000 times the amount of type I IFNs as any other cell type and both pDCs and the type I IFNs they produce have been implicated in SLE pathogenesis^{19,23}. Many genetic variants that have been identified in individuals with SLE are found in or around genes important in pDC IFN production, including *TLR7*, *SLC15A4*, *TASL*, *IRF7*, and *IRF5*^{24,25}. Type I IFNs are cytokines with myriad immune promoting functions, including promoting B cell survival, plasma cell differentiation, T cell proliferation, DC maturation as well as increasing expression of nucleic acid sensing receptors such as TLR7 and cGAS. Although these functions are critical for anti-viral immune responses, they contribute to SLE disease pathogenesis in a positive feedback loop²⁶. These type I IFNs act on many different cell types and promote the expression of interferon-stimulated genes (ISGs), many of which are

directly antiviral²⁷. ISGs are upregulated in individuals with SLE compared to healthy control individuals and correlate to measures of SLE disease severity^{28,29}. Further evidence that pDCs contribute to SLE pathogenesis comes from studies with lupus mouse models that have shown that pDC deficiency or depletion results in a reduction of lupus like-disease^{30,31}. Additionally, recent clinical trials with litifilimab, a monoclonal antibody targeting the pDC inhibitory receptor BDCA2, has shown some promising results and decreased the ISG signature in the blood of patients with SLE³². Taken together there is ample evidence that supports pDC involvement in SLE.

Plasmacytoid dendritic cells release type I IFNs in response to immune complexes

As discussed above, antinuclear antibodies are a hallmark of SLE and can form nucleic acid-containing immune complexes which deposit in tissue and can be recognized by FcR expressing cells. pDCs express the IgG specific FcγRIIa, which allows them to bind and internalize immune complexes (ICs) containing IgG¹⁵. Therefore, ANA-ICs facilitate nucleic acid entry to endosomes, where TLR7 and TLR9 reside^{15,18}. When antibodies against viral antigens deliver nucleic acids to pDC endosomes the subsequent IFN response is essential for antiviral immunity, however, during SLE, dysregulated and sustained pDC IFN production in response to self-nucleic acids contributes to pathogenesis.^{23,33}.

SLE studies have mainly focused on IgG isotype autoantibodies

While it is generally accepted that autoantibodies to nuclear antigens contribute to SLE pathogenesis there is still much to be discovered regarding how antibodies of different isotypes and antigen specificities contribute to disease. Autoantibodies of every isotype have been detected in SLE, however most studies have focused on the IgG isotype^{10,34–36}. Antibodies are immune

effector proteins produced by B cells and plasma cells. The antigen binding fragment (Fab) is the part of an antibody that interacts with antigen, confers its specificity and is unique to each B cell clone. The crystallizable fragment (Fc) is the portion of antibody that defines the five different isotypes, IgM, IgD, IgG, IgA or IgE, with each B cell clone typically only producing one isotype. IgM and IgD are the first isotypes expressed during B cell development and upon B cell receptor (BCR) recognition of antigen B cells can undergo class switching to either IgG, IgA or IgE isotype. The Fc region of each antibody isotype allows for different effector functions and recognition by corresponding receptors^{37,38}. For example, IgE immune complex recognition by FcεRI on mast cells is what drives allergic inflammation³⁹. Amounts of each antibody isotype in serum can vary greatly based on various factors including age, sex, and infection, however IgG is the most prevalent in serum with a reference range of 6.2-15.1 mg/mL, followed by IgA at 0.86-4.76 mg/mL, IgM at 0.28-2.64 mg/mL and trace amounts of IgD (~30 ug/mL) and IgE (~50 ng/mL)⁴⁰⁻⁴⁴. While most studies of autoantibodies in SLE focus on IgG, one study has looked at IgE isotype anti-dsDNA antibodies in SLE and found that they could synergize with IgG to enhance pDC responses³⁶. These findings encouraged us to investigate the role of IgA autoantibodies, especially given that there is approximately 50,000 times the amount of IgA in serum than IgE.

Evidence that IgA autoantibodies may be relevant in SLE

IgA is the second most prevalent antibody isotype in serum after IgG, and there is some evidence that IgA ANAs may contribute to SLE pathology^{35,45,46}. A few studies have shown that IgA anti-dsDNA antibodies are present in 30-50% of individuals with SLE, and that IgA anti-dsDNA antibodies correlated to SLE disease activity index (SLEDAI), lupus nephritis, and other markers of SLE severity, such as high erythrocyte sedimentation rate and low complement C3 and C4^{35,46,47}. IgA is almost uniformly present in lupus nephritis glomeruli, in addition to IgG, IgM,

C3 and C1q, and is used for diagnosis⁴⁸. Elevated levels of IgA ANAs have also been found in the saliva and fecal samples of SLE subjects^{49,50}. Additionally, SLE serum converts monocytes into dendritic cells with an acquired ability to promote both IgA and IgG plasmablast differentiation⁵¹, suggesting both IgA and IgG antibodies could contribute to disease. Lastly, one study found a substantial increase in IgA isotype B cell clones in SLE compared to healthy controls and other immune-mediated diseases⁵². However, the pathophysiologic significance of IgA in SLE is not understood and requires further investigation. Several studies have shown that IgA antibodies have a pathogenic role in other inflammatory diseases, e.g. IgA nephropathy, IgA vasculitis, dermatitis herpetiformis, inflammatory bowel disease, linear IgA bullous disease and rheumatoid arthritis⁵³.

IgA and the IgA specific receptor, Fc α R

IgA is a unique isotype that has multiple forms and functionality. IgA is generally recognized as important in mucosal immunity where it is secreted as a secretory dimeric form (SIgA) that can bind and neutralize pathogens⁵³. IgA is also abundant in human serum, where it exists mainly in a monomeric form that can be regulatory and promote immune homeostasis⁵³⁻⁵⁵. The IgA-specific Fc α R (CD89) is highly expressed in monocytes, macrophages and neutrophils. Fc α R has a minimal cytoplasmic domain and associates with the immune tyrosine activating motif (ITAM)-containing FcR common gamma (FcR γ) chain to confer downstream signaling and internalization in response to IgA-Fc α R binding^{56,57}. In monocytes, when monomeric IgA binds to Fc α R, the associated ITAMs are only partially phosphorylated resulting in recruitment of SHP-1 and downstream inhibitory signaling⁵⁸⁻⁶⁰. Multimeric or complexed IgA binding to Fc α R causes clustering of the receptor and the associated FcR γ chains resulting in complete phosphorylation of FcR γ ITAMs, recruitment of Syk, and downstream pro-inflammatory signaling^{57,61}. In this way,

IgA functions as inhibitory when monomeric and stimulatory when multimeric. In some contexts, IgA-Fc α R interactions are more effective than IgG-Fc γ R interactions at stimulating immune responses. Multimeric IgA can be more potent than IgG at activating neutrophils^{62,63} and Fc α R signaling enhances TLR signaling in monocytes and macrophages⁶⁴. Additionally, a non-synonymous SNP in *FCAR*, encoding Fc α R, which enhances signaling through this receptor, was associated with increased risk of SLE in a case-control study⁶⁵. Thus, there is evidence that IgA has more diverse roles in immune responses than previously thought and may be important in SLE. Because there is no mouse orthologue of *FCAR*, studies of IgA and Fc α R in human disease require human systems.

Results

Introduction

Because of the emerging studies suggesting IgA autoantibodies may be relevant in SLE and data demonstrating that multimeric IgA is potent at stimulating immune cells through Fc α R, we wanted to better understand the role IgA ANAs play in the context of SLE. Specifically, we asked how IgA ANAs contribute to plasmacytoid dendritic cell (pDC) responses in SLE. pDCs are specialized innate immune cells implicated in SLE pathogenesis that release large amounts of the key type I interferon (IFN) cytokines in response to RNA and DNA sensing by toll-like receptors (TLRs) 7 and 9, respectively¹⁸⁻²². Type I IFNs are cytokines critical for anti-viral immune responses with myriad immune modulating functions, including promoting B cell survival, plasma cell differentiation, T cell proliferation and DC maturation²⁶. Additionally, type I IFNs promote the expression of interferon-stimulated genes (ISGs) in many cell types, which are upregulated in individuals with SLE compared to healthy control individuals and correlate to measures of SLE disease severity^{12,23,28}. In pDCs, FcR recognition of ANA-ICs facilitates the internalization of nucleic acids to endosomes, where TLR7 and TLR9 reside^{15,18}. This leads to TLR signaling via IRF7 and subsequently potent type I IFN production, contributing to the pathogenesis of SLE^{23,33}.

Given the important role of pDCs in type I IFN production in SLE, we chose to focus on the effect IgA-containing ICs have on these cells. We found pDCs express Fc α R in addition to the IgG-binding Fc γ RIIa, and can bind both IgA and IgG. Using a unique in vitro system employing serum from individuals with SLE to generate ICs with potent pDC stimulatory function, we show here a novel and critical function for the IgA1 subtype in pDC type I IFN production in response to RNA-containing RNPs. Fc α R and Fc γ RIIa acted synergistically to induce type I IFN production

through facilitating internalization of ICs, and pDCs from individuals with SLE had higher IC internalization and Fc α R expression than those from HCs. Thus, we have uncovered a critical role of IgA1 and Fc α R in pDC responses in SLE.

SLE donors have IgA isotype ANAs to multiple nuclear autoantigens

To better understand the breadth and prevalence of these IgA anti-nuclear antibodies (ANA), we performed autoantigen profiling on serum from 24 SLE subjects assessing both IgA and IgG isotypes. Supplemental Table 1 summarizes the demographics and clinical features including disease activity scores (SLEDAI), presence of autoantibodies (ANA, dsDNA, anti-Smith RNP) and disease duration. We found that all 24 SLE donors had detectable IgA isotype autoantibodies against multiple RNA-associated and DNA-associated nuclear antigens (Fig. 1A, Supplementary Fig. 1A). When comparing IgA and IgG autoantibodies to the same autoantigen, we found that the level of these antibodies positively correlated in 25/26 nuclear antigens relevant to SLE, whereas the degree of correlation and the ratio of IgA to IgG autoantibody varied by antigen (Fig. 1A, Supplementary Fig. 1A and 1B). Antibodies against Smith/ribonucleoprotein (Sm/RNP) and double stranded DNA (dsDNA) are prevalent in SLE^{66,67} and IgA and IgG autoantibodies against these antigens were two of the most highly correlated of all nuclear antigens tested (Fig. 1B and Supplementary Fig. 1).

As an individual's serum contained both IgA and IgG autoantibodies to the same nuclear antigens, we reasoned that the immune complexes (ICs) formed in vivo when these autoantibodies recognize their cognate nuclear antigens will likely contain both IgG and IgA isotype antibodies. To test whether circulating ICs with both IgA and IgG are found in SLE, we performed ELISAs on SLE serum (n=26) using anti-IgG capture antibody and anti-IgA detection antibody. In this way we could detect ICs that contained both IgA and IgG regardless of antigen specificity. We found

that 21/26 SLE donors we tested had detectable mixed IgA/IgG ICs directly ex vivo (Fig. 1C). As has been well established⁴⁸, we detected IgA in the glomeruli of diffuse proliferative (class IV) lupus nephritis (Fig. 1D). The fact that mixed IgA/IgG ICs exist in SLE serum, that SLE serum often contains both IgA and IgG isotype ANAs, and that lupus nephritis usually shows both IgA and IgG deposition, led us to hypothesize that IgA/IgG mixed isotype ICs may contribute to disease pathogenesis in SLE. Because IgA and IgG isotype antibodies can have different effector functions and are recognized by different FcRs, these isotypes have the potential to both contribute to pathogenic IC-mediated responses in SLE.

pDCs express the IgA-specific Fc α R and bind IgA

Because pDCs make large quantities of IFN α in response to ICs, we wanted to determine how IgA-containing ICs contribute to the responses of pDCs. pDCs express the IgG-specific Fc γ RIIa (CD32a) and respond to IgG-containing ICs, but Fc α R expression on or IgA binding to pDCs has not been described. Therefore, we first examined whether pDCs express the IgA-specific Fc α R (CD89) and bind to IgA. Analysis of publicly available single-cell RNA sequencing data⁶⁸ revealed that pDCs expressed *FCAR* mRNA that encodes the IgA-specific Fc α R, albeit to a lesser degree than monocytes, which are known to have high Fc α R expression (Supplementary Fig. 2A). Interestingly, a higher percentage of pDCs had detectable *FCAR* mRNA than *FCGR1IA* mRNA encoding Fc γ RIIa, which also had lower percentage and mean counts of expression in pDCs compared to monocytes. We confirmed that pDC expression of *FCAR* mRNA resulted in Fc α R protein surface expression by flow cytometry (Fig. 2A, Supplementary Fig. 2B). We also observed Fc γ RIIa surface expression on pDCs and saw that both Fc α R and Fc γ RIIa surface expression was lower on pDCs compared to monocytes, which was consistent with the scRNA-Seq data (Fig. 2A,

2B, Supplementary Fig. 2B-C). To determine if pDCs could bind to IgA, we heat-aggregated biotinylated human IgA and used fluorescently-labeled streptavidin to visualize the IgA by flow cytometry. In this assay, pDCs bound heat aggregated IgA (Fig. 2C). In addition, we detected bound IgA on the surface of pDCs directly ex vivo by staining pDCs with anti-IgA antibody (Fig. 2D). As expected, pDCs also bound heat-aggregated IgG in vitro and had surface bound IgG directly ex vivo (Fig. 2C, 2D) Therefore, pDCs have the potential to respond to both IgA and IgG through Fc α R and Fc γ RIIIa, respectively. As we showed above that individuals with SLE have circulating ICs containing both IgA and IgG, we hypothesized that these receptors act in concert in pDCs to recognize ICs.

IgA1 in Sm/RNP immune complexes enhances pDC IFN α responses via Fc α R and Fc γ RIIIa co-engagement

We next investigated whether IgA in ICs influences pDC production of IFN α . We focused on the IgA1 subtype as this represents the majority of serum IgA^{69,70}. We modified a previously described in vitro system from *Lood et al*⁷¹ to test the effect of IgA1 autoantibodies on pDC IFN α production in response to ICs in vitro. Specifically, we used SLE serum to compare ICs that contained both IgG and IgA1 autoantibodies (SLE serum) to those depleted of IgA1 (SLE serum Δ IgA1) and those generated from purified IgA1 (SLE IgA1). (Fig. 3A). To generate these serum reagents, we first identified SLE donors in our biorepository whose serum contained both IgA and IgG specific for the RNA-containing Sm/RNP nuclear antigen. We found that 75% (12/16) of SLE donors with anti-Sm/RNP IgG also had detectable anti-Sm/RNP IgA by ELISA (Table 1, Supplementary Table 2). To generate IgA1-depleted serum (SLE serum Δ IgA1) that retained IgG and other antibody isotypes, we performed affinity chromatography on SLE serum containing anti-Sm/RNP IgA using Jacalin agarose, which binds to IgA1 (Fig. 3A). Depletion of IgA1 from serum

resulted in significant depletion of IgA anti-Sm/RNP antibodies but did not alter the amount of anti-Sm/RNP IgG or IgE as measured by ELISA (Fig. 3B, Supplementary Fig. 3A-B). IgA1-purified serum was generated by eluting the IgA1 bound to the affinity column used for generating IgA1-depleted serum. Purified IgA1 had no detectable contaminating IgG, whether specific for Sm/RNP or when measuring total IgG (Fig. 3B, Supplementary Fig. 3A-B). IgA1 depletion also did not significantly affect total or anti-Sm/RNP-specific IgE levels and purified IgA1 contained minimal anti-Sm/RNP-specific IgE (Supplementary Fig. 3A-C).

Using our defined serum reagents, we generated ICs by combining Sm/RNPs, a commercially available RNA-containing nuclear antigen, with either whole SLE serum or IgA1-depleted SLE serum to generate ICs containing both IgG and IgA or principally IgG, respectively (Fig. 3A). These ICs were added to pDCs enriched from fresh healthy control (HC) donor blood, and we measured IFN α β secretion in response to the RNA contained within the Sm/RNPs by bioassay. As expected, ICs generated by mixing SLE serum with Sm/RNPs induced a large amount of type I IFN α β secretion from HC pDCs from 4 donors, while Sm/RNPs alone elicited very low type I IFN α β production (Fig. 4A). Depletion of IgA1 from serum before generation of ICs dramatically abrogated pDC IFN α secretion, which was rescued when the purified IgA1 was added back to IgA1-depleted serum before IC generation-(Fig. 4A). Interestingly, ICs generated from purified IgA1 were not sufficient to generate strong pDC IFN α secretion, nor were ICs generated from twice the amount of IgA1-depleted serum, suggesting that IgA1 synergizes with IgG in ICs for pDC responses (Fig. 4A, Supplementary Fig. 4A).

We corroborated the importance of IgA1 in pDC responses to Sm/RNP ICs with sera from an additional 7 SLE subjects, where we found a 38-99 % reduction in IFN α secretion depending upon the serum donor (Fig. 4B, Supplementary Fig. 4B, Supplementary Table 3). We also

validated the reproducibility of these findings by repeating experiments using the same SLE serum donor with the same healthy control pDC donor over the course of 4 different experimental days months apart (Fig. 4C, Supplementary Fig. 4D). Consistent results were found when using different healthy control pDC donors with the same SLE serum donor (Supplementary Fig. 4C-D). Of note we were unable to detect any IFN β from pDCs stimulated with Sm/RNP-ICs regardless of the presence or absence of IgA1, showing that we were detecting only IFN α s in the reporter assay that measures all type I IFNs. Measurement of IFN α 2 specifically, as well as TNF, showed a similar dependence on IgA1 (Fig. 4D). Therefore, our data demonstrate a previously unreported role that IgA1 anti-nuclear antibodies have on IC activation of pDCs for cytokine secretion.

We hypothesized that by generating ICs containing both IgG and IgA when mixing SLE serum with Sm/RNPs, that co-engagement of Fc receptors for these isotypes, Fc γ RIIa and Fc α R, would be required for the potent IFN α production. To test whether Fc γ RIIa and Fc α R are required for pDC recognition of Sm/RNP ICs, we blocked pDC Fc γ RIIa with aggregated purified IgG or anti-Fc γ RIIa monoclonal antibody (mAb), or blocked Fc α R with aggregated purified IgA or anti-Fc α R mAb before addition of ICs generated with SLE serum. IC-induced IFN α was completely dependent upon signaling through Fc γ RIIa, with >95% reduction in IFN α secretion with either aggregated IgG or 6C4 blocking. Consistent with the large effect we observed of IgA1 depletion on pDC responses to ICs, both anti-Fc α R and aggregated IgA significantly reduced pDC IFN by ~80% (Fig. 4E, Supplementary Fig. 4E) These data in combination with our IgA1-depletion results led us to conclude that responses to mixed ICs containing both IgA1 and IgG are mediated through co-engagement of the IgA-specific Fc α R acting with the IgG-specific Fc γ RIIa.

IgA1 in immune complexes promotes internalization by pDCs

We hypothesized that ICs that contain IgA1 might be able to better bind to and be internalized by pDCs, delivering the Sm/RNPs to endosomes where the RNAs contained in the RNP can activate TLR7. To visualize internalization of ICs, we labeled Sm/RNPs with AF647 and generated ICs that were detectable by flow cytometry and microscopy. Using flow cytometry, we found that IC binding and internalization likely peaked ~12 hours and by 20 hours detection of Sm/RNP-AF647 decreased presumably due to degradation inside the cells (Fig. 5A). When comparing the ICs generated with IgA1-sufficient SLE serum versus those generated with IgA1-depleted SLE serum, we found that at 6 and 12 hours there was less pDC binding and internalization of IgA1-depleted ICs (Fig. 5A). In subsequent experiments we assessed the 12-hour time point and found that binding and internalization was significantly decreased in IgA1-depleted Sm/RNP ICs compared to those generated with IgA1-sufficient SLE serum using pDCs from multiple HC donors (Supplementary Fig. 5A) and using ICs generated with serum from 4 SLE donors (Fig. 5B). Therefore, these data show IgA1 promotes binding and/or internalization of Sm/RNP ICs.

When directly comparing the time course of IFN α production to corresponding Sm/RNP-AF647 internalization we found that there was detectable IFN α at 12 hours that was increased substantially at 20 hours (Supplementary Fig. 5B), while internalization peaked at 12 hours (Fig. 5A). We hypothesized that if increased internalization is the mechanism by which IgA1-IgG synergy occurs then Sm/RNP-AF647 internalization at 12 hours and IFN α production at 20 hours should correlate to one another. For this analysis we included both groups of ICs generated with either IgA1-sufficient or IgA1-depleted SLE serum. Consistent with our hypothesis, internalization of Sm/RNP ICs at 12 hours strongly correlated to IFN α secretion at 20 hours

($r=0.96$, $p=0.0002$) when assessed within a single experiment using one pDC donor and 4 SLE serum donors (Fig. 5C). When we combined data from multiple experiments significant correlation was maintained ($r=0.66$, $p=0.0005$) (Supplementary Fig. 5C) and this correlation was stronger ($r=0.79$, $p<0.0001$) when we normalized to account for the pDC donor differences in the magnitude of IFN α produced (Fig. 5D). Therefore, the degree of binding and internalization of Sm/RNP ICs correlated to the amount of IFN α produced by pDCs.

As we could not distinguish IC binding and IC internalization using the flow cytometry assay, we used confocal microscopy to assess IC internalization more directly. We added either IgA1-sufficient or IgA1-depleted AF647-labeled Sm/RNP ICs to freshly isolated HC pDCs and cultured for 12 hours, using Sm/RNP-AF647 alone as a control for non-specific uptake, and collected full z-stacks of imaged fields to determine whether the AF647 signal was fully internalized or bound to the pDC surface (Fig. 6A, Supplementary Fig. 6A, 6B). Using IgA1-sufficient SLE serum, we found that 28.5% of pDCs had internalized Sm/RNP-AF647 ICs and 8.5% had Sm/RNP-AF647 ICs that appeared bound to the surface of the cell (Fig. 6B). Sm/RNP-AF647 ICs made with SLE IgA1-depleted serum resulted in significantly reduced binding and internalization, with only 4.6% of pDCs showing internalization and 4.6% of pDCs showing binding of ICs. pDCs incubated with Sm/RNP-AF647 alone showed little binding or internalization (Fig. 6B). Taken together with our flow cytometry data, we conclude that IgA1 in ICs results in increased binding and internalization by pDCs that correlates with IFN α production. Because increased internalization would result in more effective delivery of RNA to TLR7 in the endosome, the correlation between pDC IC internalization and IFN α production is mechanistically consistent.

SLE pDCs have increased capacity to bind ICs and increased surface Fc α R expression compared to healthy control pDCs

Because IC internalization and pDC IFN α production correlated and both pDCs and IFNs are implicated in SLE pathology, we asked if pDCs from SLE donors had an increased capacity to bind to ICs compared to those from matched HC donors (Supplementary Table 3). We incubated previously frozen PBMCs from SLE and healthy control (HC) donors with Sm/RNP-AF647 ICs generated with SLE serum that contains both IgG and IgA for 3.5 hours at 37°C and then measured Sm/RNP-AF647 association with pDCs by flow cytometry (Supplementary Fig. 7A). Importantly, the SLE serum used here showed strong IgA1 dependence for pDC IFN α secretion and IC binding/internalization in our assays. We found that SLE donor pDCs had significantly increased association with Sm/RNP ICs compared to pDCs from HC donors (Fig. 7A). We also measured Fc α R and Fc γ RIIa expression on duplicate samples from this cohort and found that pDC IC binding/internalization positively correlated to pDC Fc α R surface expression, but not Fc γ RIIa surface expression (Fig. 7B). CD14⁺ monocytes in the same samples showed a trend for increased IC binding and internalization by those from SLE donors, but this did not reach statistical significance, nor did it correlate with monocyte Fc α R or Fc γ RIIa expression (Supplementary Figs. 7B, 7C). Therefore, pDCs from donors with SLE had the capacity to associate with IgA1-dependent ICs to a greater extent than those from HC donors.

Based on our findings that pDC IC binding/internalization and Fc α R expression correlated, we hypothesized that the increased Sm/RNP IC internalization by SLE pDCs might be due to increased Fc α R surface expression compared to HC pDCs, therefore we used flow cytometry to measure expression of these receptors on pDCs from 36 matched SLE and HC donors (Supplementary Table 1, 3). Fc α R expression was significantly higher on pDCs from SLE donors

than from HC donors, while there was no difference between HC and SLE pDCs in FcγRIIa expression (Fig. 7C) or IgE specific FcεR1a expression (Supplementary Fig. 7D). CD14⁺ monocytes from HC and SLE donors showed no difference in either FcαR or FcγRIIa expression (Supplementary Fig. 7E). These data suggest that the increased Sm/RNP IC binding/internalization by pDCs from SLE donors may be due to increased FcαR expression, which could lead to increased IFNα production from SLE pDCs in a positive feedback loop.

To explore whether the expression of FcαR on SLE pDCs was related to IFNα production in vivo, we performed whole blood RNA-Seq on matched samples taken from the same blood draw that we had used to measure FcαR expression for 18 subjects with samples available. Upon analysis of the top 50 genes positively correlated with FcαR MFI, we noted many were known interferon-stimulated genes (ISGs) with 19 of these top 50 found in the Hallmark IFNα response gene set (Supplementary Fig. 8A), though several others in this list are well known ISGs including *IFIT1*, *STAT1*, *OAS3* and *CCL2*. To explore this association between pDC FcαR expression and ISG signature more closely, we assessed if there was a correlation between the median expression of the entire Hallmark IFNα response gene set and pDC FcαR surface expression. There was a significant positive correlation between pDC FcαR expression in SLE donors (Fig. 7D), while there was no such correlation between pDC FcγRIIa expression or monocyte FcαR or FcγRIIa expression in these donors (Supplemental Fig. 8B-C). Therefore, the IFNα rich environment in SLE specifically correlates with FcαR expression on pDCs.

Figures:

Figure 1




Figure 1. IgA and IgG RNA-associated anti-nuclear antibodies in SLE serum.

(A) Autoantibody profiling of serum from 24 SLE subjects. Heatmaps of log-scaled antibody binding for IgA (*left*) and IgG (*right*) isotype antibodies to 16 RNA-associated nuclear antigens. Each row corresponds to an antigen and each column corresponds to an individual subject. Color bar on top indicates SLEDAI score for each donor at time of blood draw. Clustering of subjects and samples is based on similarity in scaled IgA levels, with the same clustering applied to the IgG heatmap. **(B)** Correlation between IgA and IgG for anti-Sm/RNP (*left*) and dsDNA (*right*) nuclear antigens. Each symbol represents an individual subject **(C)** Serial dilution curves of immune complexes (ICs) containing both IgA and IgG in serum from 26 SLE subjects as measured by ELISA. Of the 26 SLE subjects 15% (4/26) had no detectable IgA/IgG ICs. Each line represents an individual subject. **(D)** Histology and immunofluorescence of representative glomeruli from class IV lupus nephritis. Left Jones silver staining, and middle IgA and right IgG staining of serial sections. **(B)** r = Pearson correlation coefficient.

Figure 2. Human blood pDCs express Fc α R and Fc γ RIIa and bind IgA and IgG.

Surface staining and flow cytometry analysis performed on PBMCs from healthy control (HC) donors. **(A)** Representative histograms of Fc α R (*red*) and Fc γ RIIa (*blue*) expression on pDCs compared to mIgG1 isotype control (*grey*), with detection by the same secondary antibody. **(B)** Paired analysis between monocyte and pDC Fc α R (left) and Fc γ RIIa (right) surface staining (n=12 HC donors). **(C)** Binding of IgA and IgG to pDCs was assessed using biotinylated human heat-aggregated IgA or heat-aggregated IgG and fluorescently labeled streptavidin for visualization by flow cytometry, Streptavidin (SA) alone (*grey*), heat-aggregated IgA (*red*) and heat-aggregated IgG (*blue*). **(D)** Direct ex vivo binding of IgA and IgG to pDCs was assessed by staining with anti-IgA (*red*) or anti-IgG antibody (*blue*) directly ex vivo. Control (*grey*) has no anti-IgA or anti-IgG detection antibody. **(A-D)** Staining was performed on thawed PBMCs and pDCs gated as shown in Supplementary Fig. 2B. **(B)** Ratio paired t-test (**** p<0.0001).

Figure 3




Figure 3. RNA-containing immune complexes generated from SLE donors with anti-Sm/RNP antibodies

(A) Schematic showing reagents used to generate immune complexes (ICs) that contained both IgG and IgA (I), mainly IgG (II) or only IgA1 (III) from SLE donors with IgG and IgA anti-Sm/RNP antibodies. IgA1 depleted serum (SLE serum Δ IgA1) and purified SLE IgA1 were generated from SLE serum by using an IgA1 binding substrate column and collecting the flow through and eluted protein, respectively. ICs were made by mixing the different antibody reagents with RNA-containing Sm/RNPs. (B) Levels of IgA anti-Sm/RNP (left) and IgG anti-Sm/RNP (right) antibodies in SLE serum (purple circles), SLE serum Δ IgA1 (blue open squares) and SLE IgA1 (red diamonds), as measured by ELISA. Representative data for one serum donor is shown.

Figure 4




Figure 4. IgA1 in Sm/RNP immune complexes enhances IFN α production by human pDCs

All pDCs were enriched from PBMCs freshly isolated from healthy control (HC) donor blood. All immune complexes (ICs) were generated with Sm/RNP as the RNA-containing nuclear antigen. **(A)** pDC IFN α secretion after incubation with Sm/RNPs alone or with Sm/RNP ICs generated with SLE serum, SLE serum Δ IgA1 or SLE IgA1. Each symbol shape representing a different pDC donor, mean and range shown. **(B)** Paired analysis of pDC IFN α secretion after incubation with Sm/RNP ICs generated with either SLE serum or SLE serum Δ IgA1 from 8 SLE serum donors. **(C)** Paired analysis of pDC IFN α secretion with the same SLE serum donor and HC pDC donor assessed at 4 times over 14 months. **(D)** Paired analysis of pDC IFN α 2 (*left*) and TNF (*right*) secretion after incubation with SLE serum or SLE serum Δ IgA1 ICs. **(E)** pDC IFN α secretion after pDCs were incubated with PBS, mIgG1 isotype control, anti-Fc γ RIIa, anti-Fc α R, aggregated IgG or aggregated IgA for 2 hours prior to the addition ICs generated from SLE serum. Data are shown as a percent of PBS control with mean and range (raw data shown in Supplemental Figure 4E). **(A)** One way ANOVA, repeated measures, Bonferroni correction for multiple comparisons between all groups. Only significant comparisons are shown (*** p<0.001 and ** p<0.01). **(B-D)** Ratio paired t-test (** p<0.01 and * p<0.05). **(E)** One way ANOVA, repeated measures, Bonferroni correction for comparing all blocking conditions to the isotype control (*** p<0.001). SLE serum donors; n=1 (A, C and E), n=4 (D), n=8 (B). HC pDC donors; n=1 (B, C and D), n=3 (E), n=4 (A).

Figure 5







Figure 5: IgA1 autoantibodies contribute to IC association with pDCs

All pDCs were enriched from PBMCs freshly isolated from healthy control (HC) donor blood and ICs were generated with AlexaFluor647-labeled Sm/RNPs (Sm/RNP-AF647). **(A)** pDCs were incubated with Sm/RNP-AF647 alone (*top row*) or Sm/RNP-AF647 ICs generated from SLE serum (*middle row*) or SLE serum Δ IgA1 (*bottom row*), for 3-20 hours. Representative flow cytometry plots show the percent of Sm/RNP-AF647⁺ pDCs. The Sm/RNP-AF647⁺ gate was determined by selecting where binding of Sm/RNP-AF647 control was approximately 1% or less. **(B)** Percent IC⁺ pDCs at 12 hours after incubation with ICs generated with SLE serum or SLE serum Δ IgA1 from 4 SLE donors. **(C)** Correlation between IC⁺ pDCs (%) at 12 hrs and IFN α secretion at 20 hours (ng/mL) after incubation with Sm/RNP-AF647 ICs generated from SLE serum (*solid lavender symbols*) or SLE serum Δ IgA1 (*open blue symbols*). Plot shows the same IFN α secretion data from (B) with each symbol shape corresponding to the SLE serum donor. **(D)** Correlation of 12 hr IC⁺ pDCs (%) and normalized 20 hr IFN α production (% max) for combined experiments in which pDCs were incubated with Sm/RNP-AF647 ICs generated with SLE serum or SLE serum Δ IgA1. **(B)** Ratio paired t-test (* p<0.05) **(C, D)** r = Pearson's correlation coefficient. SLE serum donors; n=1(A), n=4 (B, C, D). HC pDC donors; n=1 (A, B, C), n=4 (D).

Figure 6




Figure 6: IgA1 autoantibodies enhance pDC internalization of immune complexes

Enriched pDCs from healthy individuals were incubated with Sm/RNP-AF647 alone or Sm/RNP-AF647 ICs generated with either SLE serum or SLE serum Δ IgA1 for 12 hrs. **(A)** Representative confocal images of the three categories of staining observed with DAPI (*blue*), CD123 (*green*) and Sm/RNP-AF647 ICs (*magenta*). Group I (*left*), pDCs that had no ICs bound or internalized or very faint IC signaling indicating low level or diffuse ICs bound. Group II (*middle*), pDCs with either small (*top cell*) or large (*bottom cell*) internalized ICs. Group III (*right*), pDCs with either small (*top cell*) or large (*bottom cell*) ICs that were bound but not internalized. Images shown are a composite of collapsed z-stacks, Supplemental Fig. 6 shows examples with full z-stacks to demonstrate the difference between internalized and bound ICs. **(B)** Donut plots show the proportion of pDCs in each group for each experimental condition: Sm/RNP-AF647 with no serum (*left*), Sm/RNP-AF647 ICs generated with SLE serum (*middle*), and Sm/RNP-AF647 ICs generated with SLE serum Δ IgA1 (*right*). The center of the donut plot shows the number of cells imaged and analyzed in each group. Chi-squared analysis performed between all groups and statistically significant comparisons shown (** $p < 0.01$, *** $p < 0.001$).

Figure 7




Figure 7. SLE pDCs have increased binding to immune complexes and increased expression of FcαR

(A) PBMCs from individuals with SLE (n=12) and matched healthy control subjects (n=12) were incubated for 3 hours with ICs generated from Sm/RNP-AF647 and SLE serum. Percent of gated pDCs with ICs is shown, determined as described in Figure 5A. (B) PBMCs from the same subjects as in (A) were also stained for surface FcαR and FcγRIIa and correlations between IC⁺ pDCs and FcαR (*left*) or FcγRIIa (*right*) are shown. Grey circles are pDCs from healthy control individuals and purple circles those from SLE individuals. (C) Cell surface expression of FcαR (*left*) and FcγRIIa (*right*) on gated pDCs from individuals with SLE (n=36) and matched HC subjects (n=37). (D) Correlation between cell surface expression of FcαRI on gated pDCs and an interferon-stimulated gene signature determined by RNA-seq performed on whole blood from 18 of the SLE subjects in (C). (A-D) Each symbol represents an individual subject; (A and C) Student's t-tests (* p<0.05 and ***p<0.001) (B and D) r = Pearson's correlation coefficient.


Tables

Table 1: SLE serum donor participants summary

SLE (n=16)	
Sex/Gender	
Female (%)	14 (87.5%)
Race	
White, Caucasian	6 (37.5%)
Black, African American	4 (25%)
Asian	2 (12.5%)
American Indian, Alaska Native	1 (6.25%)
Native Hawaiian, Other Pacific Islander	1 (6.25%)
More than one race	1 (6.25%)
Unknown	1 (6.25%)
Age at Draw	
Age at Draw (Avg +/- SD)	35.2 +/- 13.7
Age at Draw, Range	18 - 62
Clinical information	
SLEDAI Score Range	0-16
SLEDAI Score (Avg +/- SD)	4.3 +/- 5.1
SLE / Lupus Nephritis	3 (18.8%)
SLE Duration at Draw Range (yrs)	0.1-24
SLE Duration at Draw [Avg +/- SD (yrs)]	10.4 +/- 9.6
ANA Positive	16 (100%)
anti-dsDNA Positive	12 (50%)
anti-Smith, RNP or Smith RNP positive	15 (93.7%)
Experimental ELISA	
anti-Sm/RNP IgG	16 (100%)
anti-Sm/RNP IgA	12 (75%)

Supplementary Figures

Supplementary Figure 1




B

Nuclear Antigen	Mean (NSI) IgA	Mean (NSI) IgG	Ratio (NSI) IgA/IgG	(r) In IgA to IgG	(p) In IgA to IgG
CENP-A	48	915	0.05	0.13	0.5542
CENP-B	133	494	0.27	0.28	0.1781
POLB	154	836	0.18	0.39	0.0564
Genomic DNA	45	1111	0.04	0.51	0.0102
Histone H1	2855	466	6.12	0.10	0.6458
Histone H2A	4017	1189	3.38	-0.21	0.3351
Histone H2B	966	746	1.29	0.21	0.3356
Nucleosome antigen	3013	812	3.71	0.30	0.1608
dsDNA	272	3545	0.08	0.81	1.7E-06
ssDNA	1842	16628	0.11	0.88	1.4E-08
Jo-1	387	706	0.55	0.60	0.0018
La/SSB	217	643	0.34	0.66	0.0005
Ribo P. P0	462	1346	0.34	0.74	3.9E-05
Ribo P. P1	69	274	0.25	0.53	0.0082
Ribo P. P2	289	1658	0.17	0.71	3.9E-05
Ro/SSA (52 Kda)	1087	15321	0.07	0.66	0.0004
Ro/SSA (60 Kda)	110	458	0.24	0.40	0.0557
Sm	45	647	0.07	0.40	0.0545
Sm/RNP	139	2228	0.06	0.78	6.8E-06
SmD	68	672	0.10	0.52	0.0096
SmD1	143	1121	0.13	0.48	0.0167
U1-snRNP 68/70	300	5942	0.05	0.60	0.0018
U1-snRNP A	133	3272	0.04	0.54	0.0060
U1-snRNP B/B'	728	2431	0.30	0.52	0.0088
U1-snRNP C	846	3499	0.24	0.46	0.0224
ssRNA	67	299	0.22	0.25	0.2408

Supplementary Figure 1: Supplementary figures relating to main Figure 1

Autoantibody profiling of serum from 24 SLE subjects. Heatmaps of log-scaled antibody binding for IgA (*left*) and IgG (*right*) isotype antibodies to 10 DNA-associated nuclear antigens. Each row corresponds to an antigen and each column corresponds to an individual subject. The numbering of subjects below heatmap correspond to the number in Fig. 1A. The color bar across the top indicates SLEDAI score for each donor at time of blood draw. Clustering of subjects and samples is based on similarity in scaled IgA levels, with the same clustering applied to the IgG heatmap. **(B)** Table shows mean normalized signal intensity (NSI) values for IgA and IgG isotype antibody, the ratio of IgA/IgG NSI values, as well the Pearson's correlation and p values for natural log transformed IgA and IgG NSI values.

Supplementary Figure 2




Supplementary Figure 2: Supplementary figures relating to main Figure 2


(A) FCAR and FCGRIIA RNA expression in monocytes (1-4) and DC (1-6) subsets analyzed from publicly available single cell RNA sequencing data (43). Color of the circles represent scaled mean expression levels and size of the circle corresponds to the percent of expressing cells for each group. **(B)** For all flow cytometry experiments gating for pDCs was done as shown. pDCs were identified as CD3⁻CD19⁻CD14⁻CD304⁺CD123⁺ live single cells (*purple gate*). Monocyte and T cells were identified as CD3⁻CD19⁻CD14⁺ and CD19⁻CD14⁻CD3⁺ live single cells respectively. **(C)** PBMCs were analyzed by flow cytometry for surface FcαR (*red*) and FcγRIIa (*blue*) staining with isotype control staining in grey. The first two histograms show FcαR staining on gated monocytes (*first panel*) and T cells (*second panel*). The last two histograms show FcγRIIa on gated monocytes (*third panel*) and T cells (*fourth panel*). **(D)** Flow cytometry analysis showing aggregated biotinylated IgA (*red, left*) and IgG (*blue, right*) staining on monocytes as detected by labeled streptavidin. Gray histogram shows control streptavidin staining alone.

Supplementary Figure 3


A



B




C



Supplementary Figure 3: Supplementary figures relating to main Figure 3

(A) IgG (*left*), IgA (*middle*) and IgE isotype (*right*) anti-Sm/RNP was measured in arbitrary units (AU) as described in methods. SLE serum, SLE serum Δ IgA1 and purified IgA1 were assayed for the 8 SLE serum donors used for all pDC IFN α production experiments. Each donor is color coded the same throughout this figure and in all other figures. **(B)** Total IgG (*left*) in SLE serum and SLE IgA1. **(C)** Total IgA (*left*) in SLE serum and SLE serum Δ IgA1. Total IgE (*right*) in SLE serum and SLE serum Δ IgA1.


Supplementary Figure 4



Supplementary Figure 4: Supplementary figures relating to main Figure 4

All pDCs were enriched from PBMCs freshly isolated from healthy control (HC) donor blood. All immune complexes (ICs) are generated with Sm/RNP as the RNA-containing nuclear antigen **(A)** IFN α produced by pDCs after 24 incubations with Sm/RNP ICs generated with either SLE serum Δ IgA1 or twice the amount of SLE Δ IgA1 serum (2X SLE Δ IgA1 serum). **(B)** pDC IFN α production after 30-hour incubation with Sm/RNP ICs made with SLE serum or SLE serum Δ IgA1 from multiple SLE donors. Each line represents a distinct SLE serum donor. **(C)** Healthy control pDCs from multiple donors were stimulated with Sm/RNP ICs made with SLE serum or SLE serum Δ IgA1 from one SLE serum donor (same as in Fig. 4C). **(D)** (*left*) Paired analysis of pDC IFN α secretion with the same SLE serum donor and HC pDC donor assessed at 3 times over 7 months. (*right*) HC pDCs from multiple donors were stimulated with Sm/RNP ICs made with SLE serum or SLE serum Δ IgA1 from one SLE serum donor (same as *left*). **(E)** Raw non-normalized data for experiment shown in Fig. 4E. HC pDCs were blocked with either PBS, isotype control antibody, monoclonal anti-Fc γ RIIa (6C4), monoclonal anti-Fc α R (MIP8a), aggregated IgG or aggregated IgA for two hours before given Sm/RNP ICs generated with SLE serum. One way ANOVA with repeated measures ($p = 0.018$) with Bonferroni correction for multiple comparisons. Statistics for A-D are ratio paired t tests (* $p < 0.05$, ** $p < 0.01$, *** $p < 0.001$.) SLE serum donors $n=1$ (C, D & E), $n=2$ (A), $n=6$ (B *left*), $n=8$ (B *right*). HC pDC donors $n=1$ (B *left*, B *right*, D *left*), $n=3$ (A, E). $n=4$ (D *right*), $n=8$ (C).

Supplementary Figure 5




Supplementary Figure 5: Supplementary figures relating to main Figure 5


All pDCs were enriched from PBMCs freshly isolated from healthy control (HC) donor blood. All immune complexes (ICs) were generated AlexaFluor647-labeled Sm/RNPs (Sm/RNP-AF647). **(A)** Paired analysis of percent IC⁺ pDCs (n=4) after incubation with Sm/RNP-AF647 ICs generated with SLE serum or SLE serum Δ IgA1 from 1 SLE serum donor. **(B)** pDCs (n=1) were incubated with Sm/RNP-AF647 ICs generated with (n=2) SLE serum or SLE serum Δ IgA1 and IFN α secretion from replicate wells was measured at 3, 12 and 20 hours. Each graph shows data from a different SLE serum donor. **(C)** Correlation of 12 hour IC⁺ pDCs (%) and 20 hour IFN α production for combined experiments in which pDCs were incubated with ICs generated with Sm/RNP-AF647 mixed with either SLE serum or SLE serum Δ IgA1. The combined data were generated with pDCs from n=4 HC pDC donors and n=4 SLE serum donors. r = Pearson's correlation coefficient.

Supplementary Figure 6

A Sm/RNP-AF647 ICs inside pDC




B Sm/RNP-AF647 ICs outside pDC



Supplementary Figure 6: Representative images used to distinguish Sm/RNP-AF647 SLE serum immune complex internalization versus binding to pDCs

Healthy control (HC) pDCs were incubated with ICs generated with Sm/RNP-AF647 alone or Sm/RNP-AF647 with SLE serum or SLE serum Δ IgA1 for 12 hours as in described in Fig. 6. (A) Shows 3 Z stacks as a representative example of a cell with large, internalized ICs, The first image shows the middle of the cell and the last image the top of the cell. As the middle of the cell becomes more out of focus (moving left to right) the Sm/RNP-AF647 also lessens indicating that the Sm/RNP-AF647 immune complex (IC) is inside the cell. In contrast to (A), (B) shows 3 Z stacks from a representative example of a cell with large ICs bound to the outside of the cell. The first image again shows the middle of the cell and last image the top. As the middle of the cell becomes more out of focus the intensity Sm/RNP-AF647 does not change much, indicating that the Sm/RNP-AF647 IC is stuck the outside of the cell on the side. For both (A) and (B) DAPI is shown in blue, the pDC marker CD123 is shown in green and AF647 is shown in magenta.





Supplementary Figure 7



Supplementary Figure 7: Supplementary figures relating to main Figure 7

(A) Representative flow plots for data shown in Fig. 7A in which thawed and rested PBMCs from individuals with SLE (n=12) and age-, race- and sex-matched healthy control (HC) subjects (n=12) were incubated for 3 hours with ICs generated from SLE serum and Sm/RNP-AF647. The first two flow plots show the percent of Sm/RNP-AF647+ healthy control pDCs when either Sm/RNP-AF647 is added alone (first) or in combination with SLE serum (second). The second two flow plots show the percent of Sm/RNP-AF647+ SLE pDCs when either Sm/RNP-AF647 is added alone (third) or in combination with SLE serum (fourth). **(B)** Thawed and rested PBMCs from individuals with SLE (n=12) and age-, race- and sex-matched healthy control subjects (n=12) were incubated for 3 hours with ICs generated from SLE serum and Sm/RNP-AF647. Percent of gated monocytes with ICs is shown. **(C)** PBMCs from the same subjects as in **(B)** were also stained for surface Fc α R and Fc γ RIIa and correlations between IC⁺ monocytes and Fc α R (*left*) or Fc γ RIIa (*right*) staining on monocytes are shown. Grey circles are pDCs from HC individuals and purple circles those from SLE individuals. **(D)** Cell surface expression of Fc ϵ RIa on gated pDCs from individuals with SLE (n=24) and age, race, and sex-matched HC subjects (n=24). **(E)** Cell surface expression of Fc α R (*left*) and Fc γ RIIa (*right*) on gated monocytes from individuals with SLE (n=36) and age, race, and sex-matched HC subjects (n=37). **(B, D and E)** Student's t-tests **(C)** r = Pearson's correlation coefficient.

Supplementary Figure 8



Supplementary Figure 8: Top 50 genes positively correlated with Fc α R MFI from whole blood RNA-Seq

(A) Whole blood RNA-Seq was performed for n=18 donors that were also assessed for Fc α R and Fc γ RIIa expression (Fig. 7D). Heatmap shows the top 50 genes positively correlated to Fc α R expression. Scales above the heatmap show Fc α R MFI expression (red), Fc γ RIIa expression (blue) and SLEDAI score (pink). Genes that were found to be part of the Hallmark IFN-alpha response gene set from the Broad's Molecular Signatures Database are indicated with an asterisk (*). (B) Correlation between cell surface expression of Fc γ RIIa on gated pDCs and an interferon-stimulated gene signature determined by RNA-seq performed on whole blood from 18 of the SLE subjects in Fig. 7C. (C) Correlation between cell surface expression of Fc α R (*left*) and Fc γ RIIa (*right*) on gated monocytes and interferon-stimulated gene signature determined by RNA-seq performed on whole blood from 18 of the SLE subjects in Fig. 7C.

Supplementary Table 1: SLE and HC population summary

	SLE (n=24)	CONTROLS (n=25)
Sex/Gender		
Female (%)	24 (100%)	25 (100%)
Race		
White, Caucasian	16 (67%)	16 (64%)
Mixed Race	3 (12.5%)	4 (16%)
Asian	3 (12.5%)	3 (12%)
Native Hawaiian, Other Pacific Islander	0 (0%)	1 (4%)
Black, African American	2 (8%)	1 (4%)
Age at Draw		
Age at Draw (Avg +/- SD)	43.4 +/- 15.7	43.7 +/- 15.2
Age at Draw, Range	18 - 84	18 - 82
Medications		
Hydroxychloroquine	17 (68%)	n/a
Prednisone	7 (29%)	n/a
Mycophenolic Acid	3 (12.5%)	n/a
Methotrexate	2 (8%)	n/a
Azathioprine	2 (8%)	n/a
Sulfasalazine	1 (4%)	n/a
Chloroquine	1 (4%)	n/a
Clinical information		
SLEDAI Score Range	0-20	n/a
SLEDAI Score (Avg +/- SD)	4.3 +/- 4.9	n/a
SLE / Lupus Nephritis	4 (17%)	n/a
ANA Positive	24 (100%)	n/a
anti-dsDNA Positive	12 (50%)	n/a
anti-Smith positive	6 (25%)	n/a
SLE Duration at Draw Range (yrs)	0 - 22.7	n/a
SLE Duration at Draw [Avg +/- SD (yrs)]	7.4 +/- 8.5	n/a

Supplementary Table 2: SLE participant population detailed summary

SLE Serum Donor #	Sex/Gender	Age at Draw	Race	SLE: Est. Duration At Draw (Yrs)	SLEDAI Score	Tx: Azathioprine	Tx: Hydroxychloroquine	Tx: Methotrexate	Tx: Mycophenolic acid	Tx: Methylprednisone	Tx: Prednisone	Tx: Sulfasalazine	Tx: Tocilizumab	Anti Nuclear Antibody	Anti dsDNA	Anti Smith	RNP Antibodies	Smith RNP Antibodies	Serum+Sm/RNP pDC IFNg	IgG anti-Sm/RNPs	IgA anti-Sm/RNPs	Figures containing data from donor	
1	F	50	Black, African American	23.7	0						x			+	+	+	+		Y	+	+	3B, 4A-D, 5B-D, S4A, S4C, S5A, S5B(L), S5C	
2	F	62	Caucasian, American Indian, Alaska Native	17.5	0		x							+	+	+	+		Y	+	+	4B, 5B-D, 6A-B, 7A-B, S3A-B, S4D, S5B(R), S5C	
3	F	27	White, Caucasian	4.6	4		x							+	-	-	+	+	Y	+	+	4B, 5A-D, 4D-E, 6A-B, S3A-B, S4E	
4	F	51	Black, African American	11	4						x		x	+	+	+	+		Y	+	+	4B, 4D, 5B-D, S3A-B	
5	M	26	Black, African American	0.1	6	x	x				x			+	+	-	+	+	Y	+	+	4B, S3A-B	
6	F	33	White, Caucasian	15.7	9						x			+	+	+	+	+	Y	+	+	4B, S3A-B	
7	M	44	Black, African American	0.7	3	x	x							+	-	+	+	+	Y	+	+	4B, S3A-B	
8	F	39	Asian	21	0		x				x			+	+	+	+		Y	+	+	4B, 4D, S3A-B	
9	F	21	White, Caucasian	1.6	1		x	x						+	-	+	+	+	N	+	-		
10	F	57	Asian	27	4		x							+	+	+	+		N	+	+		
11	F	27	White, Caucasian	5	0		x				x			+	+	+	+		N	+	+		
12	F	22	Caucasian	9	16		x		x		x			+	+	+	+	+	N	+	+		
13	F	28	Unknown	24	14				x					+	-	-	-	+	N	+	-		
14	F	33	American Indian, Alaska Native	4	0		x		x		x			+	-	+	+		N	+	-		
15	F	24	Caucasian	2.2	0		x					x		+	-	-	-		N	+	-		
16	F	18	Native Hawaiian, Other Pacific Islander	0.2	8		x		x	x				+	-	-	+		N	+	+		
AVG		35.13		10.46	4.31	13%	75%	6%	25%	6%	50%	6%	6%	100%	56%	69%	88%	44%	50%	100%	75%		
SD		13.74		9.60	5.12																		
range		18-62		0.1-24	0-16																		

Supplementary Table 3: SLE and HC participant population summary

	SLE (n=12)	CONTROL (n=12)
Sex/Gender		
Female (%)	12 (100%)	12 (100%)
Race		
White, Caucasian	8 (66.7%)	8 (66.7%)
Asian	2 (16.7%)	2 (16.7%)
Black, African American	2 (16.7%)	2 (16.7%)
Age at Draw		
Age at Draw (Mean +/- SD)	43.8 +/- 16.1	43.8 +/-15.6
Age at Draw, Range	22-72	25-71
Medications		
Hydroxychloroquine	8 (67%)	n/a
Prednisone	2 (16.7%)	n/a
Mycophenolic Acid	2 (16.7%)	n/a
Belimumab	1 (8.3%)	n/a
Chlorquine	1 (8.3%)	n/a
Methotrexate	1 (8.3%)	n/a
Clinical information		
SLEDAI Score Range	2-9	n/a
SLEDAI Score (Avg +/- SD)	4.7 +/- 2.8	n/a
SLE / Lupus Nephritis	1 (8.3%)	n/a
ANA Positive	12 (100%)	n/a
anti-dsDNA Positive	6 (50%)	n/a
anti-Smith positive	1 (8.3%)	n/a
SLE Duration at Draw Range (yrs)	0-55	n/a
SLE Duration at Draw [Avg +/- SD (yrs)]	14.6 +/- 16.6	n/a

Supplementary Table 4: Flow cytometry antibody summary

Target	Clone	Fluorophore	Manufacturer	Catalog #	Concentration
mIgG1	RMG1-1	PE	Biologend	406608	1:50
CD3 (T cell marker)	OKT3	BV711	Biologend	317328	1:200
CD19 (B cell marker)	H1B19	FITC	Biologend	302206	1:200
CD19 (B cell marker)	H1B19	BV650	Biologend	302238	1:200
CD14 (monocyte marker)	M5E2	BUV615	BD	751150	1:200
CD123 (pDC marker)	6H6	FITC	Biologend	306014	1:50, 1:100
CD123 (pDC marker)	7G3	BUV395	BD	56419	1:50, 1:100
CD304 (pDC marker)	3E12	BV421	Biologend	145209	1:50, 1:100
FcαR	MIP8a	NA	Invitrogen	MA5-28106	1:10
FcγR1a	6C4	NA	Invitrogen	16-0329-85	1:10
isotype control mIgG1	P3.6.2.8.1	NA	Invitrogen	16-4714-85	1:10
Live/Dead	NA	NIR	Biologend	77184	1:500, 1:750

Closing Remarks

Discussion

Our study aimed to better understand how IgA isotype autoantibodies contribute to SLE pathology. Our rationale for exploring IgA isotype autoantibodies in SLE was based upon recent insights including: the presence of IgA autoantibodies in SLE ^{35,46,47,49,50,72}, evidence that IgA can be pathogenic in rheumatoid arthritis ^{73,74}, and an increasing appreciation of the ability of IgA to facilitate proinflammatory immune responses when bound to viral and bacterial pathogens or in anti-tumor responses ^{62,63}. Glomerular IgA staining in addition to IgG, IgM, C1q and C3, so-called “full house” immunofluorescence, is an almost uniform finding in all classes of lupus nephritis ⁴⁸. Additionally, IgA is the second most prevalent antibody in serum ⁵³ and is produced at the highest rate of all isotype antibodies combined, with a rate of ~66 mg/kg of body mass each day ⁵³. Using autoantigen arrays, we found that most individuals with SLE have IgA autoantibodies against at least one nuclear antigen, and these individuals often have autoantibodies of both IgA and IgG isotypes against the same nuclear antigens. We also identified circulating immune complexes containing both IgA and IgG in some individuals with SLE, suggesting that increasing our knowledge of how these isotypes work together in immune complexes is essential for better understanding the mechanisms driving SLE pathology.

We found that IgA1 autoantibodies synergize with IgG to enhance pDC responses to immune complexes in SLE. When specifically investigating anti-Sm/RNP autoantibodies, we found that most SLE donors with anti-Sm/RNP IgG also had circulating IgA to this autoantigen. Using serum from these individuals, we identified a potent synergy between IgA1 and IgG anti-Sm/RNP autoantibodies for pDC cytokine production and pDC IC internalization. Because both Fc α R and Fc γ RIIa signal through ITAMs, either in the associated FcR γ chain for Fc α R or in the

intracellular tail for Fc γ RIIa, it is not apparent why there is such strong synergy when co-signaling through these receptors when both IgG and IgA1 are present in ICs. A possible explanation for why IgA enhances the IC-mediated IFN α response is because the affinity for multimeric antigen-bound IgA for Fc α R is greater than similarly complexed IgG for Fc γ RIIa⁷⁵⁻⁷⁷. This could result in immune complexes containing both IgA1 and IgG having a higher overall avidity for pDC FcRs than those containing only IgG.

Additionally, the FcR γ signaling chain associated with Fc α R has two ITAMs, whereas Fc γ RIIa only has one ITAM⁶¹, which may lead to stronger signaling for endocytosis of mixed IgA1-IgG ICs with twice the amount of ITAM signaling per IgA1 molecule bound than per IgG molecule bound. Furthermore, ICs that contain both IgA and IgG may increase internalization by pDCs because of the increased availability of Fc receptors for binding, whereas ICs that contain one isotype of autoantibody would only be able to bind Fc α R or Fc γ RIIa, not both. The ability for an IC to bind to two different Fc receptors could be even more essential for a robust cytokine response when the surface expression of both receptors is relatively low, as is the case in pDCs, which we show express much less surface Fc α R and Fc γ RIIa than monocytes from the same individual. Interestingly, platelets are also implicated in SLE pathogenesis with reported low-level Fc α R and Fc γ RIIa expression^{78,79}. Our data showing that IC binding and Fc α R expression was greater on SLE pDCs compared to healthy controls also suggest that IgA1 enhancement of pDC IFN α production may be even more exaggerated in individuals with SLE than in healthy control individuals, exacerbating IFN α responses both to autoantigen-containing and viral-containing ICs in SLE.

IgA1 and IgG have different characteristics, which may result in qualitative changes in ICs containing both IgA1 and IgG autoantibodies compared to those without IgA1. IgA1 shares 90%

sequence identity to IgA2 with the major difference being that IgA1 has a longer hinge region with no disulfide bonds allowing for increased flexibility compared to IgA2 and IgGs^{76,80,81}. While IgG3 also has longer hinge region, it retains the disulfide bonds⁸². Interestingly one study found that longer hinge regions engineered into IgG1 and IgG3 antibodies resulted in increased phagocytosis in monocytes and neutrophils despite no change in affinity for the FcγRs⁸³. Even though this study focused on IgG hinge length and binding to FcγRs, their findings suggest that the increased hinge length of IgA1 could have a similar ability to increase internalization in pDCs via FcαR. Alternatively, the increased flexibility and length of the IgA1 hinge region may alter the size of ICs generated; larger ICs have been shown to have increased binding to FcRs^{84,85}. In addition, large ICs that contain IgA and IgG could induce a higher degree of crosslinking of FcαR and FcγRIIIa leading to stronger ITAM signaling and increased endocytosis. In our study, we did not determine the relative number or size of the ICs generated using IgA1-sufficient and IgA1-deficient SLE serum, and therefore we do not know if there were quantitative or qualitative differences in the ICs we generated. However, we found that Sm/RNP ICs generated with IgA1 alone were not sufficient to induce pDC IFNα secretion, though perhaps increasing the amounts of IgA1 used to generate ICs over the physiological amount found in SLE serum would result in pDC IFNα production with these IgA1-only ICs. The possibility remains that an excessive amount of IgA1 antibody could generate ICs capable of inducing pDC IFNα production, as IgA1 opsonized bacteria and viruses can induce phagocytosis in monocytes and phagocytosis and NETosis in neutrophils⁸⁶⁻⁸⁹, though as discussed above the low FcαR and FcγRIIIa expression on pDCs compared to these phagocytes may cause their responses to be regulated in a distinct manner. Despite these limitations, our data show a previously unrecognized connection between IgA1 and

immune complex mediated pDC IFN α production and strongly suggests that IgA1 is a critical player in SLE.

As our immune system did not evolve to cause autoimmune diseases such as SLE, we speculate that this potent synergy between IgA and IgG in ICs evolved in the context of anti-viral immune responses, for which pDC IFN α production is protective⁹⁰⁻⁹². Perhaps this synergy evolved to allow for a strong, but transitory, pDC IFN α response to virus-containing ICs. While both IgA and IgG are produced during viral infection, IgA has a much shorter half-life (~5 days) than IgG (~21 days)^{93,94}. This might allow for IgA-IgG synergy in pDC IFN α production to be important at the peak of the antibody response when viral antigens are present to generate ICs, but to resolve quickly as IgA antibodies with comparably short half-lives are removed from circulation. In support of this theory, one study quantified IgG and IgA antibodies against SARS-CoV-2 spike receptor binding domain (RBD) longitudinally after infection and found that IgA anti-RBD peaked at 16-20 days after infection and subsequently dropped to less than half the peak concentration by 41 days. In contrast, IgG anti-RBD peaked at 21-25 days and stayed relatively constant through 41 days⁹⁵. This pattern may help prevent maximal IFN α production outside the peak of the antibody response while maintaining strong lasting immunity.

In the case of SLE, self-nuclear antigens are always present at low levels making IgA-IgG synergy maladaptive in the context of autoimmunity, and may promote pathogenic, chronic IFN α production. Increased nuclear antigens due to cell death during infection or injury or due to defective clearance of dying cells as is seen often in SLE may boost nuclear antigen-specific IgA production and subsequent pDC IFN α production, promoting disease flare in SLE. Thus, determining not only the presence of nuclear antigen-specific IgA, but also how it changes over time and may fluctuate with disease flare or severity is warranted. While we did not see a

correlation between SLEDAI and IgA autoantibodies in our limited sample set, with larger cohorts there would be greater power to determine if the presence of IgA correlates to disease activity or any clinical manifestations of SLE. Additionally, because pDC Fc α R expression was higher in SLE donors and correlated with ISGs, further studies should be done to determine if pDC Fc α R expression could be a useful indicator of disease severity or activity.

The unique ability of Fc α R to transduce either activating or inhibitory signals depending on whether the bound IgA is multimeric or monomeric, respectively⁵⁸, suggests that targeting Fc α R therapeutically in SLE has two routes by which it could be beneficial. First, blocking Fc α R could potentially prevent IgA-containing immune complexes from binding to and activating Fc α R expressing cells, including pDCs, reducing IFN α production in response to ICs. If the blocking reagent is monomeric, it may also send inhibitory ITAM (ITAMi) signals through Fc α R, which have only been shown in monocytes or macrophages to date^{59,60}, but may also transduce similar signals in pDCs or neutrophils. In fact, one study has shown that early intervention targeting Fc α R with a blocking mAb resulted in reduced disease outcomes in a pristane induced lupus nephritis model in transgenic mice expressing Fc α R⁹⁶. While additional studies would be required to justify testing Fc α R as a therapeutic strategy in SLE, they are hampered by the lack of an orthologue of Fc α R in mice and therefore rely on expressing human Fc α R as a transgene using promoters that faithfully mirror expression in human.

In conclusion, we show a remarkable ability of IgA1 isotype autoantibodies to synergize with IgG in RNA-containing ICs to generate a robust pDC IFN α response in both a Fc α R and Fc γ RIIIa dependent manner. We found that this pDC IFN α response correlated to pDC IC internalization and that SLE donor pDCs exhibited higher Fc α R surface expression and more

readily internalized IgA-containing ICs compared to HC pDCs. Taken together our data indicate that IgA1 ANAs are likely contributors in SLE pathology and warrant further investigation.

Limitations

Our study has both strengths and limitations. Because we use ICs generated from SLE serum for all our studies, this has not allowed for us to define the anti-Sm/RNP IgA1 and IgG antibodies, their affinities and precise specificity, nor the optimal stoichiometry for generating pDC stimulatory ICs. We did not investigate how different IgG subtypes synergize with IgA1, nor did we isolate the effects of IgA2 subtype as most of serum IgA is IgA1 subtype^{69,70}. We also did not examine the effects of other scarce isotype antibodies, such as IgD or IgE. It is important to note that Hennault et al. found that IgG and IgE anti-dsDNA antibodies could synergize to enhance pDC IFN α responses³⁶. For this reason, we measured total IgE and anti-SmRNP IgE in our reagents and found that the effects observed were not likely to be driven by IgE as there was no significant difference in total or anti-SmRNP IgE between serum and IgA-depleted serum (Supplementary Fig. 3A and 3C). Because we did not remove IgE from our reagents it is possible that all three isotypes synergize, and future studies will focus on better defining and isolating the contributions of each isotype antibody. However, the use of our less defined SLE serum instead of monoclonal antibodies ensures that we are observing biologically disease relevant phenomenon. Additionally, isolating peripheral blood pDCs introduces donor variation and, due to the small amount of pDCs in blood, does not allow for high throughput testing of conditions optimal for inducing pDC IFN α responses to ICs. Using primary pDCs for our assays also did not allow for investigation of SLE pDC IFN α responses to ICs, as in SLE there is ~50% reduction in number of circulating pDCs and we obtain smaller amounts of blood from individuals with autoimmune

disease. However, we were able to use SLE PBMCs to investigate IC association and Fc α R expression.

Other considerations and future directions

Our findings demonstrate a novel and exciting role for IgA1 in eliciting potent IFN α production in response to RNA-containing ICs in the context of SLE, but we predict the role of IgA1-IgG synergy goes beyond Sm/RNP ICs. Thus, it is important to test if IgA1 and IgG synergy occurs when using immune complexes comprised of antibody and viral particles containing RNA to stimulate pDCs. It is also important to test whether IgA1 and IgG synergy occurs when immune complexes contain DNA thereby eliciting IFN α production via TLR9 instead of TLR7. This may be particularly interesting in the context of nucleosomes, which are important DNA-containing autoantigens in SLE. Strikingly, we found that nucleosome-specific autoantibodies, including those binding to histones H1, H2A and H2B, were the only nucleic acid associated autoantigens where there was more IgA than IgG as indicated by an IgA:IgG ratio >1 (Supplementary Fig. 1A and 1B). In preliminary experiments, we have found that SLE serum containing anti-nucleosome antibodies can form ICs with DNA-containing nucleosomes and promote pDC internalization of these ICs. Thus, we have a system in place to investigate DNA autoantigen-containing IgA1-IgG ICs to compare with SmRNP ICs.

In our studies we did not have the ability to distinguish between the different IgG subtypes (IgG1, IgG2, IgG3, and IgG4), which have different binding affinities for Fc γ RIIa⁹⁷. Therefore, we do not know if there are differences in how each individual IgG subtype synergizes with IgA1. To address this question, we will develop a system where we can generate immune complexes with known antibody isotype and subtype composition. With fluorescently-labeled Sm/RNP tetramers, we will identify B cells specific for Sm/RNPs and use single cell sorting to sequence

individual BCRs. We will then generate antibodies of each isotype and subtype (IgA1, IgA2, IgG1, IgG2, IgG3, IgG4, and IgE) with identical specificities. Using these reagents, we will generate Sm/RNP containing ICs using different antibody subtype combinations and measure pDC IFN α responses to determine the optimal IgG subtype for synergy. These reagents will also allow us to test if IgA2 synergize with IgG for pDC IC responses. Additionally, glycosylation differences have been shown to alter IgG binding to Fc γ R_s, and changes in global IgG glycosylation profiles during active disease have been described in SLE and other autoimmune diseases⁹⁸. Therefore, glycosylation states are another important variable to consider to fully understand how antibody composition influences pDC responses to immune complexes. After we successfully generate anti-Sm/RNP antibodies of each isotype/subtype we will then employ methods to test the effects of both isotype and glycosylation on pDC IFN α responses to immune complexes.

Better understanding the mechanisms by which IgA1 and IgG can synergize can help with identifying potential drug targets in SLE and provide new insights into the value of synergy during pDC immune responses to pathogens. Both Fc α R and Fc γ RIIa signal through ITAMs when immune complexes are bound, however Fc γ RIIa has one ITAM contained in its own cytoplasmic tail whereas Fc α R associates with the Fc γ chain, which contains two ITAMs. Despite both receptors using ITAMs, there must be something unique to the Fc α R cytoplasmic tail or there is some unique interaction when the Fc γ RIIa ITAM and Fc γ ITAMs are all within proximity of each other for synergy to occur when both IgA1 and IgG are present in immune complexes. Of note, there is a gain of function allele in *FCAR* that encodes an Fc α R with increased ability to recruit Lyn, a protein tyrosine kinase involved in receptor-mediated signaling⁶⁵. This allele is enriched in SLE patients of African American ancestry compared to those without SLE. Taken together, these data suggest that the cytoplasmic tail of Fc α R has some function that could result

in enhanced immune responses. Therefore, we are interested in employing various molecular techniques to interrogate the mechanisms by which Fc α R signaling synergizes with Fc γ RIIa to enhance pDC IFN responses to immune complexes.

To understand how IgA autoantibodies contribute to disease, it is important to examine IgA autoantibodies in a larger cohort of individuals with SLE. Because we show that IgA1 autoantibodies have the potential to contribute to SLE pathology in combination with IgG, we are interested in determining if the increased presence of both IgA and IgG autoantibodies correlate to disease severity, disease flare or disease manifestations. Additionally, because Fc α R expression on pDCs correlated with both IC association and ISG signature, we would like to quantify pDC Fc α R expression in a larger cohort of SLE donors to determine if pDC Fc α R expression correlated to any SLE disease measures. Determining whether Fc α R or IgA autoantibodies correlate to disease manifestations or activity is important as they both have potential as new criteria for stratification of SLE patients.

Materials and Methods

Human samples

Frozen PBMCs, frozen serum, and fresh blood draws from HC and SLE subjects were obtained from the Benaroya Research Institute Immune-mediated Disease Registry and Repository. All pDC functional assays and microscopy imaging were performed with pDCs that were magnetically enriched from PBMCs isolated on the same day as the blood draw. Frozen SLE sera was used to generate ICs and for autoantigen arrays. See supplementary tables for specifics on donors used in each experiment. All experiments comparing cells from HC and SLE subjects were performed in a blinded manner. All experiments were approved by the Benaroya Research Institute Institutional Review Board.

IgG and IgA Nuclear Antigen Microarray

Serum from 24 SLE subjects was assessed for IgG and IgA autoantigen binding at the University of Texas Southwestern (UTSW) Genomics and Microarray Core Facility. The autoantigen microarray super panel contained 128 autoantigens including most nuclear antigens relevant to SLE. For our analysis, we included 26 nuclear antigens against which SLE subjects commonly have antibodies. Normalized signal intensity (NSI) values were determined for each autoantigen as previously described by UTSW core researchers^{99,100}. These NSI values were normalized to internal controls and corresponded to the amount of antibody binding to each autoantigen regardless of isotype, which allowed for direct comparisons between IgA and IgG. NSI values were then natural log (ln) transformed for correlation analysis and antibody heatmaps were created using the BioConductor package ComplexHeatmap. Because mean NSI values varied greatly between antigens, the heatmaps shown were also scaled for each antigen so that the maximum value was “1” and minimum value was “0”. Clustering of subjects and samples was

conducted on the scaled IgA values, separately for RNA- and DNA-associated antigens, using Euclidean distances and complete linkage clustering as implemented in hclust; the clustering of IgA was then applied to the corresponding IgG heatmap. Supplementary Fig. 2B shows the raw mean NSI values.

PBMC isolation and pDC enrichment from whole blood

240 mLs of freshly drawn whole blood was diluted 1:1 with PBS and underlaid with Ficoll-Paque plus (Cytiva). After centrifugation at 1000 x g for 25 minutes at 24 C without braking, PBMCs at the interface were collected, remaining RBCs were lysed with ACK Lysis buffer (Lonza), and subsequently washed 3 times with PBS 2% FBS (Sigma). pDCs were enriched from freshly isolated PBMCs using the human Plasmacytoid Dendritic Cell Isolation Kit II (Miltenyi Biotec) following manufacturer's instructions with a 10% reduction in the recommended amounts of reagents. Enriched pDCs were counted and purity was measured by flow cytometry. pDC purity ranged from 60-99% (median 85%) with < 0.5% contamination from monocytes, B cells or T cells. For experiments using frozen PBMCs, sample vials containing 10-20 million PBMCs were thawed and washed in warmed complete RPMI (RPMI with 10% FBS, penicillin/streptomycin, L-glutamine). Complete RPMI for thawing cells also contained DNase I (Sigma) at a concentration of 9 μ g/mL.

Staining for Fc α R and Fc γ RIIa

Previously frozen PBMCs were resuspended in PBS and aliquoted into a 96 well plate for staining. All staining reagents used are listed in Supplementary Table 4. Cells were incubated with Zombie NIR live dead stain for 10 min at RT. After the live dead stain all subsequent washes and stains were performed in staining buffer (1% BSA, 2 mM EDTA in PBS) and plates were spun at

600 g for 2 min. After washing, cells were split into 3 replicate wells and incubated with purified anti-Fc α R, anti-Fc γ RIIIa or isotype control for 45 min at 4C. Samples were washed 3 times, then incubated with PE-labeled anti-mIgG1 secondary for 30 min at 4C. Samples were washed 3 times with staining buffer and once with 200 μ g/mL mouse IgG1 to bind any residual anti-mIgG1 PE. Cells were then stained with directly conjugated antibodies to CD3, CD19, CD14, CD304, and CD123 for 25 minutes at 4C, washed and fixed in 3% paraformaldehyde. For experiments where samples were stained on multiple days, a replicate vial of control donor PBMCs was stained each day and MFI values of samples were normalized to the staining of control. All flow cytometry was performed on a Cytex Aurora spectral flow cytometer. Compensation/unmixing controls were done using single color stains with antibody at the same dilution used in the complete stain. All flow cytometry analysis was done using FlowJo software.

Staining for IgA and IgG

To generate aggregated IgA or IgG, biotinylated IgG (Novus NBP1-96855) or IgA (Novus NBP1-97181) was heated at a concentration of 2 mg/mL for 35-45 min at 65 C. After live dead staining, heat aggregated antibody was diluted in staining buffer to a final concentration of 0.5 mg/mL, incubated with PBMCs for 30 min at 37C. After washing cells were incubated with PE-labeled Streptavidin followed by washing, then staining for cell specific surface markers as above. To measure IgA and IgG bound to pDCs directly ex vivo, anti-IgA (AF647) and anti-IgG (FITC) were used.

Anti-Sm/RNP and total antibody ELISAs

Flat bottom 96 well, high binding ELISA plates (Costar) were coated Sm/RNPs (Arotec Diagnostics ATR01) by adding 50 μ l Sm/RNPs (0.5-1 μ g/mL) diluted in PBS and incubated at 4

C overnight. Blocking was performed with 2X assay buffer (Invitrogen 88-50550-88) overnight at 4 C. Serum was serially diluted and incubated for 2 hrs at RT. After 7 washes with PBS/0.05% Tween-20, sample wells were incubated with biotin-labeled anti-IgA (Jackson Immunochemicals 309-065-008) or biotin-labeled anti-IgG (Jackson Immunochemicals 309-065-011) at 1:50,000 for 1 hr to detect IgA and IgG antibodies, respectively. Plates were washed as above and incubated with SA-HRP (eBiosciences 00-4100-94) for 30 mins followed by another wash cycle. TMB substrate was added, and reactions stopped with 0.16 M sulfuric acid once curves were detectable by eye (5-10 minutes). Then spectral reading was performed on VersaMax microplate reader (Molecular Devices). In addition to dilution curves, anti-Sm/RNP antigen binding was estimated in arbitrary units (AU) by using reagents from total IgG, IgA and IgE ELISA kits (ThermoFisher 88-50550-88, 88-50600-88, and 88-50610-88) to generate a standard curve for the appropriate isotype being measured using serial dilutions of IgA, IgG or IgE standards. anti-Sm/RNP isotype antibody signal was mapped on the corresponding anti-isotype antibody standard curve and scaled using arbitrary units. Total IgG, IgA and IgE were measured from serum using these same kits.

IgA/IgG IC ELISA

To detect circulating ICs from donor serum, ELISA plates were coated with anti-human IgG (ThermoFisher 88-50550-88, 1X) and blocked overnight with 2X assay buffer (Invitrogen 88-50550-88) at 4C and for 2 hours at room temperature right before use. SLE donor serum was serially diluted in assay buffer, added to wells and then incubated for 2 hours at room temperature. After 7 washes with PBS/0.05% Tween-20, biotin anti-human IgA Fab (Invitrogen A24462) was added at a dilution of 1:100,000 for 2 hours at 4 C to detect human IgA. Plates were washed again, incubated with SA-HRP for 30 mins, washing and TMB substrate detection. Reactions were

stopped with 0.16 M sulfuric acid. Spectral reading was performed on VersaMax microplate reader (Molecular Devices).

IgA1 depletion from serum

IgA1 was depleted from SLE donor serum by affinity chromatography using Jacalin agarose (Pierce catalog #20395). SLE donor serum was first diluted 1:1 in PBS and an equal volume of Jacalin agarose resin was added to the column. Resin was added to 0.8 mL or 5 mL centrifuge columns (Pierce catalog #89868 or #89896) and washed 5 times with PBS. Diluted SLE serum was added to the columns and incubated for 1 hour at room temperature. Columns were spun at 2200 x g for 5 minutes to collect IgA-depleted serum, then were washed at least 8 times or until protein A280 measurement was below 0.05 mg/mL (Nanodrop spectrophotometer ND-1000). Two washes of 1M D-Galactose (Millipore Sigma PHR1206) were used to elute IgA1 from the column. Every 1 mL of collected IgA1 was concentrated by 10X using 10K molecular weight cut off protein concentrators (Pierce 88513). After initial concentration 10 additional washes with PBS at 10X remaining sample volume were performed in the concentrators to remove any remaining D-galactose. Absorbance at 280 nm was measured to provide an estimated mg/mL amount of IgA1 solution. Aliquots were frozen at -80 C for later use.

Generation of Sm/RNP immune complexes

For each experimental well to be assayed, 12-16 μ l of SLE donor serum or IgA1-depleted SLE serum was mixed with Sm/RNPs (Arotec ATR01) at 2-3 μ g/mL in a total volume of 50 μ l (diluted in PBS) and incubated for 1 hr at room temperature. Slightly different ratios of serum and Sm/RNPs were used depending on what was optimal for any given serum donor, however ratios were kept identical for all SLE serum and Δ IgA1 serum comparisons within a given donor. In

some cases, ICs were made with Alexa Fluor 647 (AF647)-labeled Sm/RNPs. 0.5 mls of Sm/RNPs (0.75 mg/mL) were labeled using the Alexa Fluor 647 Protein Labeling Kit (Invitrogen A20173) following the manufacturer's instructions without removal of unbound AF647.

Immune complex stimulation of pDCs

25,000 – 45,000 pDCs were added to 96 well U bottom plate wells in 150 μ l of culture medium (RPMI with 2% heat inactivated human AB serum (Omega Scientific HS-20), 20 ng/mL IL-3 (Peprotech 200-03), 200 U/L penicillin and 200 μ g/L streptomycin). Cells were rested at 37°C for 1-3 hours before adding 50 μ l of Sm/RNP ICs. After the addition of the ICs, plates were incubated for 20-24 hrs then 100 μ l of supernatants were removed and stored at 4°C. For blocking experiments, 25 μ l of blocking reagents, isotype control antibody, or PBS were added to wells 2 hours before the addition of Sm/RNPs ICs. Blocking reagents were left in wells throughout the assay. Heat aggregated IgA (Invitrogen 31148) and IgG (Invitrogen UJ2861463) stock concentrations were 2 mg/mL; anti-Fc α R MIP8a (Invitrogen MA5-28106), anti-Fc γ RIIa 6C4 (Invitrogen 16-0329-85) monoclonal and isotype control (Invitrogen 16-4714-85) stock concentrations were 1 mg/mL.

Cytokine measurements from supernatants

HeLa cells stably expressing an ISRE (interferon stimulated response element) luciferase reporter construct were a generous gift from Dr. Dan Stetson (University of Washington)¹⁰¹ and were used to measure type I IFNs in supernatants. 60,000 HeLa-ISRE cells were added to each well of 96 well white opaque tissue culture plates in a volume of 100 μ l and cultured overnight. Supernatants were serially diluted in the culture medium and 50 μ l added per well. *E. coli* expressed human IFN α 1a (PBL Assay Science 11101) was serially diluted 1:2 to generate a

standard curve from 1 mg/mL to 1.9 μ g/mL. Plates were incubated at 37 C for 5-6 hours. After incubation, 50 μ l of Bright-Glo Luciferase reagent (Promega E2620) was added to each well. Luminescence was measured using a SpectraMax iD3 (Molecular Devices). In some cases, cytokines in supernatants were measured using Bioplex assays (BioLegend) measuring IFN α -2 (catalog #740351), IFN β (catalog #740353), and TNF (catalog#740359) according to manufacturer's protocols.

Interferon stimulated genes (ISGs) measurement from whole blood

Whole blood was collected in Tempus Blood RNA Tubes (ThermoFisher), and RNA was extracted using MagMax for Stabilized Blood Tubes RNA Isolation Kit, followed by globin reduction using GlobinClear Human (ThermoFisher). To generate sequencing libraries, total RNA (0.5 ng) was added to reaction buffer from the SMART-Seq v4 Ultra Low Input RNA Kit for Sequencing (Takara), and reverse transcription was performed followed by PCR amplification to generate full length amplified cDNA. Sequencing libraries were constructed using the NexteraXT DNA sample preparation kit (Illumina) to generate Illumina-compatible barcoded libraries. Libraries were pooled and quantified using a Qubit[®] Fluorometer (Life Technologies). Sequencing of pooled libraries was carried out on a NextSeq 2000 sequencer (Illumina) with paired-end 59-base reads, using a NextSeq P2 sequencing kit (Illumina) with a target depth of 5 million reads per sample. Base calls were processed to FASTQs on BaseSpace (Illumina), and a base call quality-trimming step was applied to remove low-confidence base calls from the ends of reads. The FASTQs were aligned to the GRCh38 human reference genome, using STAR v.2.4.2a, and gene counts were generated using htseq-count¹⁰². Quality metrics were calculated using the Picard family of tools (v1.134; <http://broadinstitute.github.io/picard>). We used quality thresholds of at least 2.5M total reads, at least 70% of reads aligned to human genome, and median CV of

coverage at most 0.7; 18 out of 19 samples passed these filters and were included in subsequent analyses. Gene counts were filtered to only protein-coding genes with at least one count per million in 10% of libraries and normalized using the trimmed mean of m-values¹⁰³. Differential gene expression was determined using limma-voom, with Fc α R MFI on pDCs as a continuous variable. We quantified ISG expression using the median log₂ normalized counts of gene in the MSigDB Hallmark IFN α response gene set. The gene expression heatmap was created using ComplexHeatmap.

Immune complex internalization microscopy

Freshly enriched pDCs from healthy individuals were incubated Sm/RNP-AF647 alone or with ICs generated with Sm/RNP-AF647 and SLE serum or SLE serum Δ IgA1 for 12 hrs, and then stained with anti-CD123 to identify pDCs. pDCs were mounted to glass slides using ProLongTM Diamond Antifade Mountant with DAPI (Thermofisher P36962) and imaged on Leica SP5 confocal microscope. Multiple Z-stacked images were taken per cell and images processed using Image J. pDCs were then blindly categorized into three groups (no Sm/RNP-AF647 ICs, internalized Sm/RNP-AF647 ICs or surface bound Sm/RNP-AF647 ICs) .

Human kidney biopsy

Human kidney biopsy had standard pathologic workup, including light microscopic evaluation with Jones methenamine silver, periodic acid–Schiff, hematoxylin and eosin, and trichrome stains. For immunofluorescence (IF) microscopy, frozen tissue was stained with antibodies against IgG, IgA, IgM, C3, C1q, fibrin/fibrinogen, κ light chain, lambda light chain, and albumin.

Statistics

Statistics and tests used are described in the figure legends and methods or results for each section. Calculations were performed using GraphPad Prism and R. Additionally, because Pearson correlations can be driven by outliers, we also calculated Spearman rank correlations and found qualitatively similar results.

References

1. Robinson, G. A. *et al.* Investigating sex differences in T regulatory cells from cisgender and transgender healthy individuals and patients with autoimmune inflammatory disease: a cross-sectional study. *Lancet Rheumatol.* **4**, e710–e724 (2022).
2. Roubinian, J. R., Papoian, R. & Talal, N. Androgenic Hormones Modulate Autoantibody Responses and Improve Survival in Murine Lupus. *J. Clin. Investig.* **59**, 1066–1070 (1977).
3. Taneja, V. Sex Hormones Determine Immune Response. *Front. Immunol.* **9**, 1931 (2018).
4. Souyris, M. *et al.* TLR7 escapes X chromosome inactivation in immune cells. *Sci. Immunol.* **3**, (2018).
5. Izmirly, P. M. *et al.* Prevalence of Systemic Lupus Erythematosus in the United States: Estimates From a Meta-Analysis of the Centers for Disease Control and Prevention National Lupus Registries. *Arthritis Rheumatol.* **73**, 991–996 (2021).
6. Carter, E. E., Barr, S. G. & Clarke, A. E. The global burden of SLE: prevalence, health disparities and socioeconomic impact. *Nat. Rev. Rheumatol.* **12**, 605–620 (2016).
7. Dima, A., Jurcut, C., Chasset, F., Felten, R. & Arnaud, L. Hydroxychloroquine in systemic lupus erythematosus: overview of current knowledge. *Ther. Adv. Musculoskelet. Dis.* **14**, 1759720X211073001 (2022).
8. Mackensen, A. *et al.* Anti-CD19 CAR T cell therapy for refractory systemic lupus erythematosus. *Nat. Med.* **28**, 2124–2132 (2022).
9. Murphy, G. & Isenberg, D. A. New therapies for systemic lupus erythematosus — past imperfect, future tense. *Nat. Rev. Rheumatol.* **15**, 403–412 (2019).
10. Dema, B. & Charles, N. Autoantibodies in SLE: Specificities, Isotypes and Receptors. *Antibodies* **5**, 2 (2016).
11. Ameer, M. A. *et al.* An Overview of Systemic Lupus Erythematosus (SLE) Pathogenesis, Classification, and Management. *Cureus* **14**, e30330 (2022).
12. Postal, M. *et al.* Type I interferon in the pathogenesis of systemic lupus erythematosus. *Curr. Opin. Immunol.* **67**, 87–94 (2020).
13. Renaudineau, Y., Brooks, W. & Belliere, J. Lupus Nephritis Risk Factors and Biomarkers: An Update. *Int. J. Mol. Sci.* **24**, 14526 (2023).
14. Pisetsky, D. S. & Lipsky, P. E. New insights into the role of antinuclear antibodies in systemic lupus erythematosus. *Nat. Rev. Rheumatol.* **16**, 565–579 (2020).

15. Båve, U. *et al.* FcγRIIa Is Expressed on Natural IFN-α-Producing Cells (Plasmacytoid Dendritic Cells) and Is Required for the IFN-α Production Induced by Apoptotic Cells Combined with Lupus IgG. *J. Immunol.* **171**, 3296–3302 (2003).
16. Guilliams, M., Bruhns, P., Saeys, Y., Hammad, H. & Lambrecht, B. N. The function of Fcγ receptors in dendritic cells and macrophages. *Nat. Rev. Immunol.* **14**, 94–108 (2014).
17. Wang, Y. & Jönsson, F. Expression, Role, and Regulation of Neutrophil Fcγ Receptors. *Front. Immunol.* **10**, 1958 (2019).
18. Means, T. K. *et al.* Human lupus autoantibody–DNA complexes activate DCs through cooperation of CD32 and TLR9. *J Clin Invest* **115**, 407–417 (2005).
19. Siegal, F. P. *et al.* The Nature of the Principal Type 1 Interferon-Producing Cells in Human Blood. *Science* **284**, 1835–1837 (1999).
20. Swiecki, M. & Colonna, M. The multifaceted biology of plasmacytoid dendritic cells. *Nat. Rev. Immunol.* **15**, 471–485 (2015).
21. Panda, S. K., Kolbeck, R. & Sanjuan, M. A. Plasmacytoid dendritic cells in autoimmunity. *Curr. Opin. Immunol.* **44**, 20–25 (2017).
22. Reizis, B. Plasmacytoid Dendritic Cells: Development, Regulation, and Function. *Immunity* **50**, 37–50 (2019).
23. Elkon, K. B. & Stone, V. V. Type I Interferon and Systemic Lupus Erythematosus. *J. Interf. Cytokine Res.* **31**, 803–812 (2011).
24. Brown, G. J. *et al.* TLR7 gain-of-function genetic variation causes human lupus. *Nature* **605**, 349–356 (2022).
25. Iwamoto, T. & Niewold, T. B. Genetics of human lupus nephritis. *Clin. Immunol.* **185**, 32–39 (2017).
26. McNab, F., Mayer-Barber, K., Sher, A., Wack, A. & O’Garra, A. Type I interferons in infectious disease. *Nat. Rev. Immunol.* **15**, 87–103 (2015).
27. Schoggins, J. W. & Rice, C. M. Interferon-stimulated genes and their antiviral effector functions. *Curr. Opin. Virol.* **1**, 519–525 (2011).
28. Baechler, E. C. *et al.* Interferon-inducible gene expression signature in peripheral blood cells of patients with severe lupus. *Proc. Natl. Acad. Sci.* **100**, 2610–2615 (2003).
29. Nehar-Belaid, D. *et al.* Mapping systemic lupus erythematosus heterogeneity at the single-cell level. *Nat. Immunol.* **21**, 1094–1106 (2020).

30. Rowland, S. L. *et al.* Early, transient depletion of plasmacytoid dendritic cells ameliorates autoimmunity in a lupus model. *J. Exp. Med.* **211**, 1977–1991 (2014).
31. Sisirak, V. *et al.* Genetic evidence for the role of plasmacytoid dendritic cells in systemic lupus erythematosus. *J. Exp. Med.* **211**, 1969–1976 (2014).
32. Furie, R. *et al.* Monoclonal antibody targeting BDCA2 ameliorates skin lesions in systemic lupus erythematosus. *J. Clin. Investig.* **129**, 1359–1371 (2019).
33. Honda, K. & Taniguchi, T. IRFs: master regulators of signalling by Toll-like receptors and cytosolic pattern-recognition receptors. *Nat. Rev. Immunol.* **6**, 644–658 (2006).
34. Vladutiu, A. O. Immunoglobulin D: Properties, Measurement, and Clinical Relevance. *Clin. Diagn. Lab. Immunol.* **7**, 131–140 (2000).
35. Villalta, D. *et al.* Anti-dsDNA Antibody Isotypes in Systemic Lupus Erythematosus: IgA in Addition to IgG Anti-dsDNA Help to Identify Glomerulonephritis and Active Disease. *PLoS ONE* **8**, e71458 (2013).
36. Henault, J. *et al.* Self-reactive IgE exacerbates interferon responses associated with autoimmunity. *Nat. Immunol.* **17**, 196–203 (2016).
37. James, L. K. B cells defined by immunoglobulin isotypes. *Clin. Exp. Immunol.* **210**, 230–239 (2022).
38. Chiu, M. L., Goulet, D. R., Teplyakov, A. & Gilliland, G. L. Antibody Structure and Function: The Basis for Engineering Therapeutics. *Antibodies* **8**, 55 (2019).
39. Rios, E. J. & Kalesnikoff, J. Mast Cells, Methods and Protocols. *Methods Mol. Biol.* **1220**, 239–255 (2014).
40. Laurent, J. *et al.* [How to define the normal level of serum IgE in adults?]. *Ann. Med. interne* **136**, 419–22 (1985).
41. Gonzalez-Quintela, A. *et al.* Serum levels of immunoglobulins (IgG, IgA, IgM) in a general adult population and their relationship with alcohol consumption, smoking and common metabolic abnormalities. *Clin. Exp. Immunol.* **151**, 42–50 (2008).
42. Khan, S. R. *et al.* Determinants and Reference Ranges of Serum Immunoglobulins in Middle-Aged and Elderly Individuals: a Population-Based Study. *J. Clin. Immunol.* **41**, 1902–1914 (2021).
43. Rogentine, G. N., Rowe, D. S., Bradley, J., Waldmann, T. A. & Fahey, J. L. Metabolism of human immunoglobulin D (IgD). *J. Clin. Investig.* **45**, 1467–1478 (1966).

44. Kelly, B. T. & Grayson, M. H. Immunoglobulin E, what is it good for? *Ann. Allergy, Asthma Immunol.* **116**, 183–187 (2016).
45. Lee, A. Y. S. IgA anti-dsDNA antibodies: A neglected serological parameter in systemic lupus erythematosus. *Lupus* **31**, 137–142 (2021).
46. Jost, S. A. *et al.* IgG, IgM, and IgA Antinuclear Antibodies in Discoid and Systemic Lupus Erythematosus Patients. *Sci. World J.* **2014**, 171028 (2014).
47. Witte, T. *et al.* Association of IgA anti-dsDNA antibodies with vasculitis and disease activity in systemic lupus erythematosus. *Rheumatol. Int.* **18**, 63–69 (1998).
48. Weening, J. J. *et al.* The classification of glomerulonephritis in systemic lupus erythematosus revisited. *Kidney Int.* **65**, 521–530 (2004).
49. Romero-Ramírez, S. *et al.* Salivary IgA subtypes as novel disease biomarkers in systemic lupus erythematosus. *Front. Immunol.* **14**, 1080154 (2023).
50. Gudi, R., Kamen, D. & Vasu, C. Fecal immunoglobulin A (IgA) and its subclasses in systemic lupus erythematosus patients are nuclear antigen reactive and this feature correlates with gut permeability marker levels. *Clin. Immunol.* **242**, 109107 (2022).
51. Joo, H. *et al.* Serum from patients with SLE instructs monocytes to promote IgG and IgA plasmablast differentiation. *J. Exp. Med.* **209**, 1335–1348 (2012).
52. Bashford-Rogers, R. J. M. *et al.* Analysis of the B cell receptor repertoire in six immune-mediated diseases. *Nature* **574**, 122–126 (2019).
53. Gool, M. M. J. van & Egmond, M. van. IgA and Fc α RI: Versatile Players in Homeostasis, Infection, and Autoimmunity. *ImmunoTargets Ther.* **9**, 351–372 (2021).
54. Olas, K., Butterweck, H., Teschner, W., Schwarz, H. P. & Reipert, B. Immunomodulatory properties of human serum immunoglobulin A: anti-inflammatory and pro-inflammatory activities in human monocytes and peripheral blood mononuclear cells. *Clin. Exp. Immunol.* **140**, 478–490 (2005).
55. Wolf, H. M. *et al.* Human serum IgA downregulates the release of inflammatory cytokines (tumor necrosis factor- α , interleukin-6) in human monocytes. *Blood* **83**, 1278–88 (1994).
56. Morton, H. C. *et al.* Functional Association between the Human Myeloid Immunoglobulin A Fc Receptor (CD89) and FcR γ Chain. *J. Biol. Chem.* **270**, 29781–29787 (1995).
57. Shen, L. *et al.* Presentation of ovalbumin internalized via the immunoglobulin-A Fc receptor is enhanced through Fc receptor γ -chain signaling. *Blood* **97**, 205–213 (2001).

58. Pasquier, B. *et al.* Identification of Fc α RI as an Inhibitory Receptor that Controls Inflammation Dual Role of FcR γ ITAM. *Immunity* **22**, 31–42 (2005).
59. Kanamaru, Y. *et al.* Inhibitory ITAM Signaling by Fc α RI-FcR γ Chain Controls Multiple Activating Responses and Prevents Renal Inflammation. *J. Immunol.* **180**, 2669–2678 (2008).
60. Watanabe, T. *et al.* Negative regulation of inflammatory responses by immunoglobulin A receptor (Fc α RI) inhibits the development of Toll-like receptor-9 signalling-accelerated glomerulonephritis. *Clin. Exp. Immunol.* **166**, 235–250 (2011).
61. Mkaddem, S. B., Benhamou, M. & Monteiro, R. C. Understanding Fc Receptor Involvement in Inflammatory Diseases: From Mechanisms to New Therapeutic Tools. *Front. Immunol.* **10**, 811 (2019).
62. Brandsma, A. M. *et al.* Potent Fc Receptor Signaling by IgA Leads to Superior Killing of Cancer Cells by Neutrophils Compared to IgG. *Front. Immunol.* **10**, 704 (2019).
63. Gimpel, A.-K. *et al.* IgA Complexes Induce Neutrophil Extracellular Trap Formation More Potently Than IgG Complexes. *Front. Immunol.* **12**, 761816 (2022).
64. Hansen, I. S., Hoepel, W., Zaat, S. A. J., Baeten, D. L. P. & Dunnen, J. den. Serum IgA Immune Complexes Promote Proinflammatory Cytokine Production by Human Macrophages, Monocytes, and Kupffer Cells through Fc α RI–TLR Cross-Talk. *J. Immunol.* **199**, 4124–4131 (2017).
65. Wu, J. *et al.* Fc α RI (CD89) Alleles Determine the Proinflammatory Potential of Serum IgA. *J. Immunol.* **178**, 3973–3982 (2007).
66. Ahn, S. S. *et al.* Anti-Smith antibody is associated with disease activity in patients with new-onset systemic lupus erythematosus. *Rheumatol. Int.* **39**, 1937–1944 (2019).
67. Rekvig, O. P. Anti-dsDNA antibodies: Critical remarks. *Clin. Exp. Immunol.* **179**, 5–10 (2015).
68. Villani, A.-C. *et al.* Single-cell RNA-seq reveals new types of human blood dendritic cells, monocytes, and progenitors. *Science* **356**, (2017).
69. Delacroix, D. L. *et al.* Changes in size, subclass, and metabolic properties of serum immunoglobulin A in liver diseases and in other diseases with high serum immunoglobulin A. *J. Clin. Investig.* **71**, 358–367 (1983).
70. Brown, T. A., Murphy, B. R., Radl, J., Haaijman, J. J. & Mestecky, J. Subclass distribution and molecular form of immunoglobulin A hemagglutinin antibodies in sera and nasal secretions after experimental secondary infection with influenza A virus in humans. *J. Clin. Microbiol.* **22**, 259–264 (1985).

71. Lood, C. *et al.* C1q inhibits immune complex–induced interferon- α production in plasmacytoid dendritic cells: A novel link between C1q deficiency and systemic lupus erythematosus pathogenesis. *Arthritis Rheum.* **60**, 3081–3090 (2009).
72. Ukadike, K. C. *et al.* IgG and IgA autoantibodies against L1 ORF1p expressed in granulocytes correlate with granulocyte consumption and disease activity in pediatric systemic lupus erythematosus. *Arthritis Res. Ther.* **23**, 153 (2021).
73. Aleyd, E., Al, M., Tuk, C. W., Laken, C. J. van der & Egmond, M. van. IgA Complexes in Plasma and Synovial Fluid of Patients with Rheumatoid Arthritis Induce Neutrophil Extracellular Traps via Fc α RI. *J. Immunol.* **197**, 4552–4559 (2016).
74. Rantapää-Dahlqvist, S. *et al.* Antibodies against cyclic citrullinated peptide and IgA rheumatoid factor predict the development of rheumatoid arthritis. *Arthritis Rheum.* **48**, 2741–2749 (2003).
75. Bruhns, P. *et al.* Specificity and affinity of human Fc γ receptors and their polymorphic variants for human IgG subclasses. *Blood* **113**, 3716–3725 (2009).
76. Cottignies-Calamarte, A., Tudor, D. & Bomsel, M. Antibody Fc-chimerism and effector functions: When IgG takes advantage of IgA. *Front. Immunol.* **14**, 1037033 (2023).
77. Bakema, J. E. & Egmond, M. van. The human immunoglobulin A Fc receptor Fc α RI: a multifaceted regulator of mucosal immunity. *Mucosal Immunol.* **4**, 612–624 (2011).
78. Qian, K. *et al.* Functional expression of IgA receptor Fc α RI on human platelets. *J. Leukoc. Biol.* **84**, 1492–1500 (2008).
79. Linge, P., Fortin, P. R., Lood, C., Bengtsson, A. A. & Boilard, E. The non-haemostatic role of platelets in systemic lupus erythematosus. *Nat. Rev. Rheumatol.* **14**, 195–213 (2018).
80. Woof, J. M. & Russell, M. W. Structure and function relationships in IgA. *Mucosal Immunol.* **4**, 590–597 (2011).
81. Sousa-Pereira, P. de & Woof, J. M. IgA: Structure, Function, and Developability. *Antibodies* **8**, 57 (2019).
82. Liu, H. & May, K. Disulfide bond structures of IgG molecules. *mAbs* **4**, 17–23 (2012).
83. Chu, T. H. *et al.* Hinge length contributes to the phagocytic activity of HIV-specific IgG1 and IgG3 antibodies. *PLoS Pathog.* **16**, e1008083 (2020).
84. Lux, A., Yu, X., Scanlan, C. N. & Nimmerjahn, F. Impact of Immune Complex Size and Glycosylation on IgG Binding to Human Fc γ Rs. *J. Immunol.* **190**, 4315–4323 (2013).

85. Chen, H. *et al.* Detection and functional resolution of soluble immune complexes by an Fc γ R reporter cell panel. *EMBO Mol. Med.* **14**, e14182 (2022).
86. Stacey, H. D. *et al.* IgA potentiates NETosis in response to viral infection. *Proc. Natl. Acad. Sci.* **118**, e2101497118 (2021).
87. Aleyd, E. *et al.* IgA Enhances NETosis and Release of Neutrophil Extracellular Traps by Polymorphonuclear Cells via Fc α Receptor I. *J. Immunol.* **192**, 2374–2383 (2014).
88. Duchemin, M., Tudor, D., Cottignies-Calamarte, A. & Bomsel, M. Antibody-Dependent Cellular Phagocytosis of HIV-1-Infected Cells Is Efficiently Triggered by IgA Targeting HIV-1 Envelope Subunit gp41. *Front. Immunol.* **11**, 1141 (2020).
89. Davis, S. K., Selva, K. J., Kent, S. J. & Chung, A. W. Serum IgA Fc effector functions in infectious disease and cancer. *Immunol. Cell Biol.* **98**, 276–286 (2020).
90. Cella, M., Facchetti, F., Lanzavecchia, A. & Colonna, M. Plasmacytoid dendritic cells activated by influenza virus and CD40L drive a potent TH1 polarization. *Nat. Immunol.* **1**, 305–310 (2000).
91. Swiecki, M., Gilfillan, S., Vermi, W., Wang, Y. & Colonna, M. Plasmacytoid Dendritic Cell Ablation Impacts Early Interferon Responses and Antiviral NK and CD8⁺ T Cell Accrual. *Immunity* **33**, 955–966 (2010).
92. Yun, T. J. *et al.* Human plasmacytoid dendritic cells mount a distinct antiviral response to virus-infected cells. *Sci. Immunol.* **6**, (2021).
93. Morell, A., Skvaril, F., Nosedá, G. & Barandun, S. Metabolic properties of human IgA subclasses. *Clin. Exp. Immunol.* **13**, 521–8 (1973).
94. Morell, A., Terry, W. D. & Waldmann, T. A. Metabolic properties of IgG subclasses in man. *J. Clin. Investig.* **49**, 673–680 (1970).
95. Zeng, W. *et al.* Characterization of SARS-CoV-2-specific antibodies in COVID-19 patients reveals highly potent neutralizing IgA. *Signal Transduct. Target. Ther.* **6**, 35 (2021).
96. Liu, C. *et al.* Targeted IgA Fc receptor I (Fc α RI) therapy in the early intervention and treatment of pristane-induced lupus nephritis in mice. *Clin. Exp. Immunol.* **181**, 407–416 (2015).
97. Li, X., Ptacek, T. S., Brown, E. E. & Edberg, J. C. Fc γ receptors: structure, function and role as genetic risk factors in SLE. *Genes Immun.* **10**, 380–389 (2009).
98. Zhou, X. *et al.* Antibody glycosylation in autoimmune diseases. *Autoimmun. Rev.* **20**, 102804 (2021).

99. Robinson, W. H. *et al.* Autoantigen microarrays for multiplex characterization of autoantibody responses. *Nat. Med.* **8**, 295–301 (2002).
100. Li, Q. -Z. *et al.* Protein array autoantibody profiles for insights into systemic lupus erythematosus and incomplete lupus syndromes. *Clin. Exp. Immunol.* **147**, 60–70 (2007).
101. Lau, L., Gray, E. E., Brunette, R. L. & Stetson, D. B. DNA tumor virus oncogenes antagonize the cGAS-STING DNA-sensing pathway. *Science* **350**, 568–571 (2015).
102. Anders, S., Pyl, P. T. & Huber, W. HTSeq—a Python framework to work with high-throughput sequencing data. *Bioinformatics* **31**, 166–169 (2015).
103. Robinson, M. D. & Oshlack, A. A scaling normalization method for differential expression analysis of RNA-seq data. *Genome Biol.* **11**, R25 (2010).