


# UNIVERSITY OF WASHINGTON DEPARTMENT OF OCEANOGRAPHY

Technical Reports  
Nos. 142, 143, 144, 145, 146,  
147, 148, and 149

A COMPILATION OF ARTICLES REPORTING RESEARCH  
Sponsored Jointly by  
THE U.S. ATOMIC ENERGY COMMISSION  
and  
THE OFFICE OF NAVAL RESEARCH

U.S. Atomic Energy Commission  
Contract AT(45-1)-1725  
and  
Office of Naval Research  
Contracts Nonr-477(10)  
and Nonr-477(37)  
Project NR 083 012

Reference M66-2  
January 1966



SEATTLE, WASHINGTON 98105

UNIVERSITY OF WASHINGTON  
DEPARTMENT OF OCEANOGRAPHY  
Seattle, Washington 98105

Technical Reports

Nos. 142, 143, 144, 145, 146,  
147, 148, and 149

A COMPILATION OF ARTICLES REPORTING RESEARCH


SPONSORED JOINTLY BY

THE U.S. ATOMIC ENERGY COMMISSION


and

THE OFFICE OF NAVAL RESEARCH

U. S. Atomic Energy Commission  
Contract AT(45-1)-1725  
and  
Office of Naval Research  
Contracts Nonr-477(10)  
and Nonr-477(37)  
Project NR 083 012

  
RICHARD H. FLEMING  
Chairman

Reference M66-2  
January 1966

  
CLIFFORD A. BARNES  
Principal Investigator

Reproduction in whole or in part is permitted  
for any purpose of the United States Government

ARTICLES REPORTING RESEARCH SPONSORED JOINTLY BY THE  
U.S. ATOMIC ENERGY COMMISSION AND THE OFFICE OF NAVAL RESEARCH

## Technical Report No. 142

BIO-LITHOLOGY OF NORTHEAST PACIFIC SURFACE SEDIMENTS, by Y. R. Nayudu and B. J. Enbysk. *Marine Geology*, 2(4):310-342. 1964. (AEC: RLO-1725-42)

## Technical Report No. 143

THE TINTINNID PARAFAVELLA GIGANTEA (BRANDT), KOFOID & CAMPBELL, 1929, IN THE NORTH PACIFIC OCEAN, by Hsin-Yi Ling. *Journal of Paleontology*, 39(4):721-723. 1965. (AEC: RLO-1725-43)

## Technical Report No. 144

THE CARBONATE CONTENT OF SURFACE SEDIMENTS FROM THE NORTHEAST PACIFIC OCEAN, by M. Grant Gross. *Northwest Science*, 39(3):85-92. 1965. (AEC: RLO-1725-44)

## Technical Report No. 145

RADIOACTIVITY OF THE COLUMBIA RIVER EFFLUENT, by M. Grant Gross, Clifford A. Barnes, and Gordon K. Riel. *Science*, 149(3688):1088-1090. 1965. (AEC: RLO-1725-45)

## Technical Report No. 146

FRACTIONATION OF PHYTOPLANKTON COMMUNITIES OFF THE WASHINGTON AND OREGON COASTS, by George C. Anderson. *Limnology and Oceanography*, 10(3):477-480. 1965. (AEC: RLO-1725-46)

## Technical Report No. 147

CHLOROPHYLLS IN MARINE PHYTOPLANKTON: CORRELATION WITH CARBON UPTAKE, by G. C. Anderson and K. Banse. *Deep-Sea Research*, 12(4):531-533. 1965. (AEC: RLO-1725-47)

## Technical Report No. 148

GRAPHIC REPRESENTATION OF THE SALINITY DISTRIBUTION NEAR THE COLUMBIA RIVER MOUTH, by Betty-Ann Morse and Noel McGary. Pp. 923-942 in *Ocean Science and Ocean Engineering* 1965, vol. 2. Marine Technology Society, Washington, D. C. 1965. (AEC: RLO-1725-3)

## Technical Report No. 149

THE UNION OF THE COLUMBIA RIVER AND THE PACIFIC OCEAN -- GENERAL FEATURES, by Alyn C. Duxbury. Pp. 914-922 in *Ocean Science and Ocean Engineering* 1965, vol. 2. Marine Technology Society, Washington D. C. 1965. (AEC: RLO -1725-5)

## FRACTIONATION OF PHYTOPLANKTON COMMUNITIES OFF THE WASHINGTON AND OREGON COASTS<sup>1</sup>

Fractionation of phytoplankton communities into different size groups has been carried out in several previous studies. The seasonal changes in the distribution of various size groups have been measured in inshore marine waters (Yentsch and Ryther 1959; Holmes and Anderson 1963) and in lakes (Rodhe, Vollenweider, and Nauwerck 1958). Saijo (1964) reported on the size distribution of phytoplankton in the Indian Ocean at different depths and areas. All of these studies have emphasized the importance of the con-

tribution of the nanoplankton to the total photosynthesis of phytoplankton communities. The present study shows the seasonal and annual changes in the size distribution of phytoplankton in areas of the northeast Pacific Ocean ranging from coastal waters to the open ocean.

Fractionation of surface phytoplankton communities was carried out routinely as part of a continuing oceanographic investigation concerning the effects of the Columbia River effluent on the northeast Pacific Ocean. The information is pertinent to food chain considerations and to studies of the turnover of radionuclides introduced into the Columbia River by operations of the Hanford Laboratories. During 25 cruises from 1961-1963, 307 separate fractionations were made. Daily, at a station taken near noon, a 500-ml sample of sur-

<sup>1</sup> Contribution No. 348 from the Department of Oceanography, University of Washington. The assistance of Miss Ann Helander is gratefully acknowledged. Financial support was provided by U.S. Atomic Energy Commission Contract No. AT(45-1)-1725 and Office of Naval Research Contract Nonr-477 (10), Project NR 083 012 and Contract Nonr-477 (37), Project ~~RR-004-03-01-NR 053 012~~




FIG. 1. Index map showing the area of investigation along the Washington-Oregon coast. The shaded area indicates the combined coverage of all cruises.

face water was inoculated with approximately  $5 \mu\text{C}$   $^{14}\text{C}$  as  $\text{Na}_2\text{CO}_3$  and incubated under artificial illumination (9,150 lux) at sea surface temperature. The  $^{14}\text{C}$ -labeled population was fractionated by passing aliquots through a series of filters (nylon net- $35.0 \mu$ ; Millipore<sup>®2</sup> filters: SM- $5.0 \mu$ , AA- $0.8 \mu$ , HA- $0.45 \mu$ , PH- $0.3 \mu$ , GS- $0.22 \mu$ ) with a vacuum of 375-500 Torr. Radioactivity was measured with a gas flow counter with a micromil window. The mean value of the HA, PH, and GS filters was taken as 100% retention because each of these filters retained about the same activity when less than 500 Torr of vacuum was used.

The data are grouped into three areas, continental shelf, continental slope, and oceanic waters (Fig. 1), because it is known that the species composition is quite different in

coastal and in oceanic areas (Hobson 1964) and that phytoplankton population size and productivity are generally much greater in the coastal area (Anderson 1964). In the oceanic area, phytoplankton productivity is characterized by a spring bloom and a lesser autumn pulse, but the standing crop of phytoplankton as measured by chlorophyll *a* shows little seasonal variation with the exception of a distinct summer minimum. In coastal areas, both phytoplankton productivity and standing stock are sustained at high levels throughout the summer because of an abundant nutrient supply from upwelled waters.

The variations in retention by each filter pore size were erratic but presumably reflect the changes in size composition of the community (Fig. 2, Table 1). As would be expected, coarser filters retained less activity than finer filters. The retention by the  $35\text{-}\mu$  net was small in most cases but ranged from 2 to 39% retention. Usually rather large fractions of the phytoplankton passed through the  $5\text{-}\mu$  filter (42-100% retention), whereas the amount passing through the  $0.8\text{-}\mu$  filter was generally small (76-100% retention). In view of the large variations in short intervals of time and lack of apparent trends, annual or seasonal mean values for retention by filters of pore size greater than  $0.45 \mu$  have little significance and therefore will not be reported.

There were no regular seasonal trends exhibited by any cell-size group nor were there any great annual or geographic differences. Over the continental shelf, there was a greater preponderance of larger organisms ( $> 35 \mu$ ) during the summer months. The water over the continental slope also showed a slight trend towards larger organisms during summer, but this trend in oceanic waters was not evident. However, the oceanic area apparently had, at most times, more fragile cells or possibly smaller cells than did coastal areas, especially in 1961 and 1963, as evidenced by the amount of material passed by the  $0.8\text{-}\mu$  filter.

The material passing through the  $5\text{-}\mu$  filter might be expected to be the small

<sup>2</sup> Registered trademark, Millipore Filter Corporation, Bedford, Massachusetts.




FIG. 2. The relative retention of  $^{14}\text{C}$ -labeled phytoplankton by a graded series of filters from areas off the Washington-Oregon coast during 1961-1963. The number above each histogram designates the number of separate fractionations made during each cruise. The width of the histogram is proportional to the duration of the cruise. \*—35- and 0.45- $\mu$  filters only were used during this cruise.

microflagellates and possibly small elongate diatoms that happened to be oriented in the proper direction. Because all phytoplankton cells are larger than 1  $\mu$ , as far as is known, phytoplankton material that passes a 0.8- $\mu$  filter would apparently be

TABLE 1. The range of relative retention (%) of  $^{14}\text{C}$ -labeled phytoplankton by a graded series of filters from areas off the Washington-Oregon coast during 1961-1963

		Continental shelf	Continental slope	Oceanic
35 $\mu$	1961	4-23	2-13	2-12
	1962	4-30	3-16	3-29
	1963	3-28	3-39	11-31
5 $\mu$	1961	70-87	70-99	70-82
	1962	81-98	76-92	76-100
	1963	58-100	42-97	63-87
0.8 $\mu$	1961	84-100	77-100	83-90
	1962	86-98	76-92	89-100
	1963	76-100	88-99	81-100

either fragments of cells or possibly small pliable cells that have been squeezed through the pores by suction pressure. Direct evidence for the existence of whole cells in the filtrate is not available, but significant photosynthetic activity has been measured in samples inoculated with  $^{14}\text{C}$  after filtration through a 0.8- $\mu$  filter. In four experiments carried out in 1960 at the Friday Harbor Laboratories,  $^{14}\text{C}$  uptake measurements in seawater samples previously filtered through 0.8- $\mu$  filters ranged from 1.2 to 6.3% of total  $^{14}\text{C}$  uptake in raw seawater samples after correction for dark uptake. However, considerable fragmentation of cells could have taken place during filtration because in these experiments a vacuum as high as 685 Torr was used.

Microscopic examination of the phytoplankton during January 1961, showed the microflagellates to be an important part of

the community and were as much as three times greater in biomass than the diatom population in some areas (Hobson 1964). The significant loss of material through the 5- $\mu$  filter at that time suggests that the difference between the SM (5.0  $\mu$ ) and HA (0.45  $\mu$ ) filters gives some idea of the abundance of microflagellates in the community, although the estimate is not as great as that shown by microscopic examination. A large fraction of particles smaller than Millipore filter pore size is retained by secondary valence forces of the filters (Millipore Filter Corp. 1964) and as a result would lead to minimal estimates for the smaller particles. Because it is not known what fraction of particles is held by secondary valence forces, fractionation with Millipore filters is not quantitative. However, the amount of phytoplankton activity generally passed by the 5- $\mu$  and 0.8- $\mu$  filters suggests that microflagellates at most times may be much more important in the ocean than has been previously thought.

GEORGE C. ANDERSON

*University of Washington,  
Department of Oceanography,  
Seattle.*

#### REFERENCES

- ANDERSON, G. C. 1964. The seasonal and geographic distribution of primary productivity off the Washington and Oregon coasts. *Limnol. Oceanog.*, **9**: 284-302.
- HOBSON, L. A. 1964. Some influences of the Columbia River effluent on marine phytoplankton during January, 1961. Dept. of Oceanography, Univ. Washington, Tech. Rept. No. 100, (Ref. M64-18). 46 p.
- HOLMES, R. W., AND G. C. ANDERSON. 1963. Size fractionation of C<sup>14</sup>-labeled natural phytoplankton communities, p. 241-250. *In* C. H. Oppenheimer [ed.], *Symposium on marine microbiology*. Thomas, Springfield, Ill.
- MILLIPORE FILTER CORPORATION. 1964. Millipore ® filters and associated apparatus. Catalog MF-64. Bedford, Massachusetts. 48 p.
- RODHE, W., R. A. VOLLENWEIDER, AND A. NAUWERCK. 1958. The primary production and standing crop of phytoplankton, p. 299-322. *In* A. A. Buzzati-Traverso [ed.], *Perspectives in marine biology*. University of California.
- SAIJO, Y. 1964. Size distribution of photosynthesizing phytoplankton in the Indian Ocean. *J. Oceanog. Soc. Japan*, **19**: 19-21.
- YENTSCH, C. A., AND J. H. RYTHER. 1959. Relative significance of the net phytoplankton and nannoplankton in the waters of Vineyard Sound. *J. Conseil, Conseil Perm. Intern. Exploration Mer*, **24**: 231-238.